
On the social cost of carbon and discounting in the DICE model

Philipp Braun^{1,*}, Timm Faulwasser², Lars Grüne³, Christopher M. Kellett¹, Willi Semmler⁴, and Steven R. Weller⁵,

¹ School of Engineering, The Australian National University, Canberra, Australia

² Institute of Energy Systems, Energy Efficiency and Energy Economics, Technische Universität Dortmund, Dortmund, Germany

³ Mathematisches Institut, Universität Bayreuth, Bayreuth, Germany

⁴ New School for Social Research, New York, USA, and Center for European Economic Research, Germany, and International Institute for Applied Systems Analysis, Austria

⁵ School of Engineering, University of Newcastle, Newcastle, Australia

* **Correspondence:** philipp.braun@anu.edu.au.

Abstract: The social cost of carbon (SCC) has emerged as one of the relevant measures in integrated assessment models in climate economy, to quantify costs related to global warming and climate change. While the SCC is used in different models including DICE (Dynamic Integrated model of Climate and Economy), PAGE (Policy Analysis of the Greenhouse Effect) and FUND (Climate Framework for Uncertainty, Negotiation, and Distribution), its exact definition and computation depends on the reference and frequently lacks consistency within research streams focusing on a single or on different models. In this paper, we investigate three different methods for the computation of the SCC using the integrated assessment model DICE. While the first two methods are commonly known and used, the novel formula derived for the third method allows a direct analysis of the impact of the discount factor in the calculation of the SCC. We provide a detailed proof for the correctness of the third method and validate all three methods by numerical experiments.

Keywords: economics of climate change; DICE; optimal control; social cost of carbon; sensitivity analysis

1. Introduction

To quantify the damages from anthropogenic emissions of heat-trapping greenhouse gases, economists model the dynamics of climate–economy interactions using integrated assessment models (IAMs), which incorporate mathematical models of phenomena from both geophysical science and economics. A central role for IAMs is to estimate the social cost of carbon (SCC), defined as the

dollar value of the economic damage caused by a one metric tonne increase in carbon dioxide (CO₂) emissions to the atmosphere. The SCC can then form the basis of a Pigouvian tax on greenhouse gas emissions, guide carbon prices in emissions trading schemes, inform cost-benefit analyses driving climate policy, or be used by investors looking to consider potential emissions costings in investment decisions. This wide range of uses highlights the importance of methods for computing the SCC.

A critical and much debated issue in computing the SCC is the choice of discount rate. CO₂ is a long-lived greenhouse gas (GHG), with CO₂ emissions having a substantial impact on climate for thousands of years; e.g. 20–40% of a CO₂ perturbation remains in the atmosphere even after equilibration with the ocean on a timescale of 2–20 centuries [1, 2]. Comparing costs and benefits over such long time horizons necessitates a careful choice of discount rates and methodologies in order to provide a fair comparison of costs paid now—via emissions reductions—with economic benefits from curbed global warming in the far distant future.

Conceptually, the most common approach to computing the SCC using IAMs is to compute baseline emissions and economic consumption. These computations are then repeated with an increment of CO₂ emissions in a chosen year, yielding reduced economic consumption into the future as the result of a climate damages function in the IAM. The SCC is computed as the net present value (NPV) of the “damage stream” (i.e. pathway of lost consumption) (see, for example, [3, 4]). This NPV approach is employed directly by two of the most widely used IAMs: PAGE (Policy Analysis of the Greenhouse Effect [5]) and FUND (Climate Framework for Uncertainty, Negotiation, and Distribution [6]). While descriptively this is what is intended with DICE (Dynamic Integrated model of Climate and Economy, see Figure 1 and associated discussion in [4]), in practice the publicly available implementation of DICE uses a different method, based on shadow prices or Lagrange multipliers obtained as a result of solving an optimal control problem (see [7]).

The classical Ramsey formula for social discount rate is given by $r = \rho + \theta g$, where ρ is the pure rate of time preference (or “impatience factor”), θ is the elasticity of marginal utility of consumption, and g is the growth rate of per-capita consumption [8]. While the calculation of the SCC as the NPV of damage stream due to CO₂ pulse implies use of a discount rate, there is widespread disagreement in the literature regarding the appropriate discount rate to use in this context; see for example the following list of representative papers, wherein the SCC is described as being computed as the NPV of a damage stream variously discounted at: (a) the classical Ramsey formula (i.e., assuming a constant growth rate of consumption); (b) a time-varying Ramsey formula (reflecting the growth rate of time-varying consumption); (c) the pure rate of time preference; and (d) a fixed rate without any explicit reference to the Ramsey formula; see [9], [6], [10], [11], [5], [12], [13], [3], [14], [15, Appendix E], [4], [16], and also [17, Section 13.7] which proposes exercises based on both (b) and (d). The report [18, Chapter 7] provides a chapter on various discounting approaches in the economics of climate change. It is worth noting that in some cases it is unclear precisely how the discounting is being performed as it is described in words rather than by providing an explicit mathematical expression.

As briefly mentioned above, a conceptually distinct approach to computing SCC relies on maximizing a social welfare function. As explicitly defined in [19], “The [SCC] in a particular year is the decrease in aggregate consumption in that year that would change the current expected value of social welfare by the same amount as a one unit increase in carbon emissions in that year.” Expressed

formally, the SCC at time t is defined as

$$S_{cc}(t) = -\frac{\frac{\partial W}{\partial E(t)}}{\frac{\partial W}{\partial C(t)}} \quad (1.1)$$

where a simple manipulation shows that this is the ratio of the marginal prices of consumption and emissions. In fact, this is the computation contained in the code implementing the DICE model and optimal control problem as presented in [7]. Here, W denotes social welfare, E denotes emissions and C denotes consumption, whose definitions are made precise in Section 2.

The above discussion indicates three possible computational methods for the SCC: a direct discretization of (1.1); using the ratio of marginals obtained from solving an optimal control problem; and an NPV approach based on experimental CO₂ emission pulses. Additionally, for the last approach, there are several options for how to perform the required discounting. This leads us to the following question: which (if any) of the above options for computing the SCC are consistent with each other? Importantly, neither FUND nor PAGE involve an optimal control problem, meaning the first two methods mentioned here are not relevant. Hence, answering this consistency question is critical for comparing results between IAMs based on optimal control methodologies with those based on scenario methodologies.

Our main mathematical result is Theorem 1, which demonstrates that in order to obtain consistency between the NPV calculation and the SCC values calculated via optimal control approaches, it is necessary to use a time-varying discount rate. In particular, the time-varying component is inversely proportional to economic growth, leading to a declining discount rate [20], [21], [22]. Specifically, returning to the Ramsey formula $r = \rho + \theta g$, mentioned above, the quantity g is time-varying and given by the growth rate of consumption. The quantities ρ and θ are “exogenous” to the model and need to be selected by the modeler on the basis of data and/or economic expertise. Note that it has been argued that the use of declining rates is justified in the presence of significant uncertainty in future economic growth (see [23]). Here, we see that the use of a declining discount rate is required for consistency with outputs from calculations based on shadow prices obtained when solving optimal control problems.

This paper is organized as follows. In Section 2 we succinctly present the DICE model as described in [24]. In Section 3, we investigate three methods for computation of the SCC using the DICE model. Section 3.1 presents the calculation based on a discretization of the difference quotient of the two quantities in (1.1). In Section 3.2 we demonstrate how to rewrite the DICE optimal control problem so that emissions and consumption are decision variables, which allows us to obtain the marginals or dual variables directly as an output of solving the optimal control problem. This has the computational benefit of removing the welfare function as an intermediary in the SCC calculation. In Section 3.3 we present the main mathematical result of the paper (Theorem 1) which shows an equivalence between calculating the SCC as a ratio of the dual variables versus the NPV when the discounting approach is computed via the time-varying Ramsey formula accounting for a time-varying growth rate of consumption. Section 4 presents some numerical results and discussion and we conclude in Section 5. Mathematical necessities and the proof of our main result are collected in Appendices.

Notation: The set $\mathbb{N}_0 = \{0, 1, 2, \dots\}$ denotes the natural numbers including zero, while $\mathbb{N} = \{1, 2, \dots\}$ denotes the natural numbers without zero. The real numbers are denoted as \mathbb{R} and $\mathbb{R}_{\geq 0} = \{x \in \mathbb{R} \mid x \geq 0\}$ denotes the real numbers bigger or equal to zero. For $n \in \mathbb{N}$, the set $\mathbb{R}^n = \mathbb{R} \times \dots \times \mathbb{R}$ denotes the n -fold

Cartesian product of the real numbers and $\boldsymbol{x} \in \mathbb{R}^n$ denotes an n -dimensional vector with $x(i) \in \mathbb{R}$, $i \in \{1, \dots, n\}$. To simplify expressions and to define convergence rates we use the big O -notation, i.e., two functions $f_1, f_2 : \mathbb{R} \rightarrow \mathbb{R}$, $f_2(x) > 0$ for all $x > 0$, satisfy $f_1(x) = O(f_2(x))$ if and only if there are $C > 0$ and $\delta > 0$ such that $|f_1(x)| \leq C f_2(x)$ for all $0 < x < \delta$. For a continuously differentiable function $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$, $(x, y) \mapsto f(x, y)$, the gradient is denoted by $\nabla f(x, y)$. The gradient and the second derivative of f with respect to a subset of variables, that is with respect to x , for example, is denoted by $\nabla_x f(x, y)$ and $\nabla_{xx}^2 f(x, y)$, respectively.

2. The DICE model

For the computation of the social cost of carbon, we consider the Dynamic Integrated model of Climate and Economy (DICE). For a comprehensive introduction of the model and its parameters we refer to [4, 24]. Here we only present the equations of the model that are necessary for the following sections while following the notation introduced in [24]. We focus on the parameters defining the R-DICE2016 model, but the results derived in this paper also hold for the parameters defining the DICE-2013R model. Following the R-DICE2016 setting we denote time in years by $t \in \mathbb{R}_{\geq 0}$ so that

$$t = t_0 + \Delta \cdot i, \quad (2.1)$$

with initial time $t_0 = 2015$, a sample period $\Delta = 5$, and $i \in \mathbb{N}_0$. The DICE model is described through the dynamics

$$M(i+1) = \Phi_M M(i) + B_M \left(\sigma(i)(1 - \mu(i))A(i)K(i)^\gamma L(i)^{1-\gamma} + E_{\text{Land}}(i) + E_{\text{Pulse}}(i) \right), \quad (\text{CAR})$$

$$T(i+1) = \Phi_T T(i) + B_T \left(F_{2 \times} \log_2 \left(\frac{M_{\text{AT}}(i)}{M_{\text{AT},1750}} \right) + F_{\text{EX}}(i) \right), \quad (\text{CLI})$$

$$K(i+1) = \Phi_K K(i) + \Delta \left(\frac{1}{1 + a_2 T_{\text{AT}}(i)^{a_3}} \right) \left(1 - \theta_1(i)\mu(i)^{\theta_2} \right) A(i)K(i)^\gamma L(i)^{1-\gamma} s(i), \quad (\text{CAP})$$

$$L(i+1) = L(i) \left(\frac{1 + L_a}{1 + L(i)} \right)^{\ell_g}, \quad (\text{POP})$$

$$A(i+1) = \frac{A(i)}{1 - g_A \exp(-\delta_A \Delta(i-1))}, \quad (\text{TFP})$$

$$\sigma(i+1) = \sigma(i) \exp\left(-g_\sigma(1 - \delta_\sigma)^{\Delta(i-1)} \Delta\right), \quad (\text{EI})$$

modeling the climate (CLI), the carbon cycle (CAR), the capital (CAP), the population (POP), the total factor productivity (TFP), and the emissions intensity (EI). The remaining time dependent quantities are described through the equations

$$\theta_1(i) = \frac{P_b}{1000 \cdot \theta_2} (1 - \delta_{pb})^{i-1} \cdot \sigma(i), \quad (2.2)$$

$$F_{\text{EX}}(i) = f_0 + \min \left\{ f_1 - f_0, \frac{f_1 - f_0}{t_f} (i-1) \right\}, \quad (2.3)$$

$$E_{\text{Land}}(i) = E_{L0} \cdot (1 - \delta_{EL})^{i-1}. \quad (2.4)$$

Control inputs are the savings rate, $s(i) \in [0, 1]$ for all $i \in \mathbb{N}_0$, and the mitigation rate, $\mu(i) \in [0, 1]$ for all $i \in \mathbb{N}_0$. An additional parameter, $E_{\text{Pulse}}(i) \in \mathbb{R}$, for all $i \in \mathbb{N}_0$, is used for the calculation of the partial

derivatives needed contained in the definition of the SCC, cf. equation (3.2), below. The additional parameter $E_{\text{Pulse}}(i)$ has an impact on the total emissions

$$E(i) = \sigma(i)(1 - \mu(i))A(i)K(i)^\gamma L(i)^{1-\gamma} + E_{\text{Land}}(i) + E_{\text{Pulse}}(i) \quad (2.5)$$

at time $i \in \mathbb{N}$, which we can identify in (CAR). The parameter is not present in the baseline DICE model where $E_{\text{Pulse}}(i) = 0$ for all $i \in \mathbb{N}_0$. We collect the states in the variable X , i.e.,

$$X(i) = \{T(i), M(i), K(i), L(i), A(i), \sigma(i)\}$$

and use $X(0) = X_0$ to denote a fixed initial condition.

The control inputs are used to maximize the social welfare depending on the consumption and the utility at time $i \in \mathbb{N}_0$ as

$$C(i) = \left(\frac{1}{1 + a T_{\text{AT}}(i)^2} \right) \left(1 - \theta_1(i)\mu(i)^{\theta_2} \right) A(i)K(i)^\gamma L(i)^{1-\gamma} (1 - s(i)) + C_{\text{Pulse}}(i) \quad (2.6)$$

$$U(i) = L(i) \left(\frac{\left(\frac{1000C(i)}{L(i)} \right)^{1-\alpha} - 1}{1 - \alpha} \right). \quad (2.7)$$

Here, $C_{\text{Pulse}}(i) \in \mathbb{R}$, for all $i \in \mathbb{N}_0$, plays the same role as $E_{\text{Pulse}}(i)$ in Equation (CAR), manipulating the consumption in a specific year. Then, for a fixed horizon $N \in \mathbb{N}$, the social welfare is defined as

$$W := \sum_{i=0}^{N-1} \frac{U(i)}{(1 + \rho)^{\Delta i}} \quad (2.8)$$

and the optimal control problem (2.9) which maximizes the social welfare and generates optimal pathways is defined as

$$\begin{aligned} W^* &= \max_{s, \mu} \sum_{i=0}^{N-1} \frac{U(i)}{(1 + \rho)^{\Delta i}} \\ &\text{subject to } X(0) - X_0 = 0 \\ &\quad (\text{CAR}) - (\text{EI}), \quad \forall i = 0, \dots, N-1, \\ &\quad (2.2) - (2.4), \quad \forall i = 0, \dots, N-1, \\ &\quad s(i), \mu(i) \in [0, 1] \quad \forall i = 0, \dots, N-1. \end{aligned} \quad (2.9)$$

The optimal control problem (2.9) constitutes a nonlinear program (NLP). We use the notation W^* to denote the optimal value with respect to $C_{\text{Pulse}}(j) = E_{\text{Pulse}}(j) = 0$, for all $j \in \{0, \dots, N-1\}$, i.e., W^* denotes the nominal or baseline solution of (2.9). In the case that $C_{\text{Pulse}}(j) \neq 0$ or $E_{\text{Pulse}}(j) \neq 0$ for $j \in \{0, \dots, N-1\}$, is considered in (2.9), the optimal solution is denoted by $W^*[C_{\text{Pulse}}(j)]$ or $W^*[E_{\text{Pulse}}(j)]$, respectively. Similarly, optimal pathways are denoted using \cdot^* . In this context, $C^*(\cdot)$, $C^*[C_{\text{Pulse}}(j)](\cdot)$, and $C^*[E_{\text{Pulse}}(j)](\cdot)$ denote the optimal consumption pathways obtained through (2.9) with $C_{\text{Pulse}}(j) = E_{\text{Pulse}}(j) = 0$, $C_{\text{Pulse}}(j) \neq 0$, and $E_{\text{Pulse}}(j) \neq 0$, respectively (in all cases we set $C_{\text{Pulse}}(i) = E_{\text{Pulse}}(i) = 0$ for all $i \neq j$). Additionally, $N \in \mathbb{N}$ denotes a fixed horizon and $s, \mu \in \mathbb{R}^N$ define a short-hand notation for

$$s = \begin{bmatrix} s(0) \\ \vdots \\ s(N-1) \end{bmatrix} \quad \text{and} \quad \mu = \begin{bmatrix} \mu(0) \\ \vdots \\ \mu(N-1) \end{bmatrix}. \quad (2.10)$$

3. Calculation of the Social Cost of Carbon

In this section we take as a formal definition of the SCC at time $j \in \{0, \dots, N-1\}$, a scaled version of the change of the optimal consumption $C^*(j)$ with respect to a change of the optimal emissions $E^*(j)$, i.e., we define

$$S_{CC}(j) = -1000 \frac{\partial C^*(j)}{\partial E^*(j)}. \quad (3.1)$$

The scaling provides an SCC value in US dollars per tonne of CO₂, [24, Sec. 2.8]. Since $E^*(j)$ is itself an outcome of the optimal control problem (2.9) rather than an independent variable on which $C^*(j)$ would depend, it is not immediately clear how this partial derivative has to be interpreted. Its meaning is as follows: we consider $E_{\text{Pulse}}(j) \in \mathbb{R} \setminus \{0\}$ and $E_{\text{Pulse}}(i) := 0$ for all $i \neq j$. Then we solve the optimal control problem (2.9) in order to obtain $C^*[E_{\text{Pulse}}(j)](\cdot)$. The partial derivative in (3.1) is then defined as

$$\frac{\partial C^*(j)}{\partial E^*(j)} := \lim_{\substack{E_{\text{Pulse}}(j) \rightarrow 0 \\ E_{\text{Pulse}}(j) \neq 0}} \frac{C^*[E_{\text{Pulse}}(j)](j) - C^*(j)}{E_{\text{Pulse}}(j)}. \quad (3.2)$$

In the same way, partial derivatives with respect to other quantities occurring in the optimization can be defined.

Under appropriate differentiability assumptions on E^* , C^* and the optimal value function W^* , Equation (3.1) can be rewritten as

$$S_{CC}(j) = -1000 \frac{\partial C^*(j)}{\partial W^*} \cdot \frac{\partial W^*}{\partial E^*(j)} = -1000 \frac{\partial W^* / \partial E^*(j)}{\partial W^* / \partial C^*(j)} \quad (3.3)$$

used as the common definition in [4, Sec. 2.1], for example. Here, “[t]he numerator is the marginal impact of emissions at time j on welfare, while the denominator is the marginal welfare value of a unit of aggregate consumption in period j ”, [4, Sec. 2.1].

Since the optimal value function W^* as well as the optimal total emissions $E^*(j)$ and optimal consumptions $C^*(j)$ are only implicitly known as the optimal solutions of the optimal control problems (2.9), a direct calculation of the social cost of carbon through (3.3) is not possible. Nevertheless, there are several efficient ways to approximate (3.3).

3.1. Direct discretization based on the difference quotient

Using (3.2), we can approximate the quantity in (3.3) at a fixed time $j \in \{0, \dots, N-1\}$ by performing three experiments to compute approximations

$$\frac{\partial W^*}{\partial E^*(j)} \approx \frac{W^*[E_{\text{Pulse}}(j)] - W^*}{E_{\text{Pulse}}(j)} \quad \text{and} \quad \frac{\partial W^*}{\partial C^*(j)} \approx \frac{W^*[C_{\text{Pulse}}(j)] - W^*}{C_{\text{Pulse}}(j)}. \quad (3.4)$$

To this end three optimal control problems can be solved.

1. A baseline solution of (2.9) that generates a welfare value W^* corresponding to $E_{\text{Pulse}}(j) = 0$ and $C_{\text{Pulse}}(j) = 0$ for all $j = 0, \dots, N-1$.
2. A second solution of (2.9) with a small non-zero perturbation on emissions at time $j \in \{0, \dots, N-1\}$; i.e., $E_{\text{Pulse}}(j) \neq 0$. This yields a second (optimal) welfare value $W^*[E_{\text{Pulse}}(j)]$.

3. A third solution of (2.9) with a small non-zero perturbation on consumption at time $j \in \{0, \dots, N-1\}$; i.e., $C_{\text{Pulse}}(j) \neq 0$ that yields a third (optimal) welfare value $W^*[C_{\text{Pulse}}(j)]$.

Then the SCC in (3.3) can be approximated by

$$S_{\text{CC}}(j) \approx -1000 \frac{C_{\text{Pulse}}(j)}{E_{\text{Pulse}}(j)} \cdot \frac{W^*[E_{\text{Pulse}}(j)] - W^*}{W^*[C_{\text{Pulse}}(j)] - W^*}. \quad (3.5)$$

Since (3.4) is a first order approximation of the derivative of W^* with respect to $E^*(j)$ and $C^*(j)$, respectively, the right-hand side of (3.5) converges with order $O(C_{\text{Pulse}}(j) + E_{\text{Pulse}}(j))$ to (3.3) for $C_{\text{Pulse}}(j) \rightarrow 0$ and $E_{\text{Pulse}}(j) \rightarrow 0$ and under the assumption that the optimal control problem (2.9) can be solved with arbitrary precision. To calculate $S_{\text{CC}}(j)$ for all $j \in \{0, \dots, N-1\}$ using the approximation (3.5), $2N+1$ optimal control problems need to be solved.

Remark 1. *The order of convergence follows from Taylor expansions of W^* under the assumption that W^* is twice continuously differentiable with respect to E and C . In particular, the derivation of the error term $O(C_{\text{Pulse}}(j) + E_{\text{Pulse}}(j))$ relies on*

$$\begin{aligned} W^*[E_{\text{Pulse}}(j)] &= W^* + \frac{\partial W^*}{\partial E^*(j)} E_{\text{Pulse}}(j) + O(E_{\text{Pulse}}(j)^2) \\ W^*[C_{\text{Pulse}}(j)] &= W^* + \frac{\partial W^*}{\partial C^*(j)} C_{\text{Pulse}}(j) + O(C_{\text{Pulse}}(j)^2) \end{aligned} \quad (3.6)$$

and is given in detail in Appendix C.

3.2. SCC calculation based on dual variables

Most modern NLP solvers, including IPOPT [25] and CONOPT [26], for example, the optimal Lagrange multipliers, i.e., the dual variables, of all explicitly enforced equality and inequality constraints in addition to the optimal solution W^* , s^* , μ^* . This can be used to compute the SCC directly via dual variables. All what is necessary to do is formulating (2.9) in a way such that $C(i)$ and $E(i)$ are decision variables of the NLP. Using (2.5) allows a reformulation of (CAR) in terms of

$$M(i+1) = \Phi_M M(i) + B_M E(i). \quad (3.7)$$

Instead of (2.9), we can thus consider the modified optimal control problem

$$\begin{aligned} W^* &= \max_{s, \mu, C, E} \sum_{i=0}^{N-1} \frac{U(i)}{(1+\rho)^{\Delta i}} \\ \text{subject to } & X(0) - X_0 = 0 \\ & (3.7), (\text{CLI}) - (\text{EI}), \quad \forall i = 0, \dots, N-1, \\ & (2.2) - (2.4), \quad \forall i = 0, \dots, N-1, \\ & (2.5) - (2.6), \quad \forall i = 0, \dots, N-1, \\ & s(i), \mu(i) \in [0, 1] \quad \forall i = 0, \dots, N-1. \end{aligned} \quad (3.8)$$

where

$$C = \begin{bmatrix} C(0) \\ \vdots \\ C(N-1) \end{bmatrix} \quad \text{and} \quad E = \begin{bmatrix} E(0) \\ \vdots \\ E(N-1) \end{bmatrix}$$

are defined similar to (2.10). The optimal control problems (2.9) and (3.8) are identical in the sense that the optimal solutions W^* as well s^* and μ^* coincide. However, in (3.8) we avoid the substitution of (2.6) and (2.5).

The main difference between (2.9) and (3.8) is that in the latter case, consumption and emissions are treated as *pseudo* decision variables defined via explicit equality constraints, while in the former case one substitutes $C(i)$ and $E(i)$ in the objective and the dynamics by the expressions given in (2.6) and (2.5). The implementation of (3.8) in an NLP solver such as IPOPT yields values for the optimal Lagrange multipliers of all constraints.

To make this more precise, note that we can rewrite (3.8) in form of the parameter dependent optimization problem

$$\begin{aligned} f^*(\mathbf{p}) = \max_{\mathbf{x}} f(\mathbf{x}) \\ \text{subject to } \quad & h_{E_i}(\mathbf{x}) = p_{E_i}, \quad \forall i = 0, \dots, N-1, \\ & h_{C_i}(\mathbf{x}) = p_{C_i}, \quad \forall i = 0, \dots, N-1, \\ & k(\mathbf{x}) = 0, \\ & g(\mathbf{x}) \geq 0, \end{aligned} \quad (3.9)$$

in the optimization variables $\mathbf{x} = \{s, \mu, \mathbf{C}, \mathbf{E}\}$. Here,

$$f(\mathbf{x}) = \sum_{i=0}^{N-1} \frac{U(i)}{(1+\rho)^{\Delta i}}$$

denotes the objective function in the variables $s, \mu, \mathbf{C}, \mathbf{E}$ and

$$\begin{aligned} h_{E_i}(\mathbf{x}) &= E(i) - (\sigma(i)(1 - \mu(i))A(i)K(i)^\gamma L(i)^{1-\gamma} + E_{\text{Land}}(i)), \\ h_{C_i}(\mathbf{x}) &= C(i) - \left(\left(\frac{1}{1 + a T_{\text{AT}}(i)^2} \right) (1 - \theta_1(i)\mu(i)^{\theta_2}) A(i)K(i)^\gamma L(i)^{1-\gamma} (1 - s(i)) \right), \end{aligned}$$

capture (2.5) and (2.6) for $i = 0, \dots, N-1$. The pulses E_{Pulse} and C_{Pulse} are included through the parameter $\mathbf{p} = [p_E^T \ p_C^T]^T$,

$$\mathbf{p}_E = \begin{bmatrix} p_{E_0} \\ \vdots \\ p_{E_{N-1}} \end{bmatrix} \quad \text{and} \quad \mathbf{p}_C = \begin{bmatrix} p_{C_0} \\ \vdots \\ p_{C_{N-1}} \end{bmatrix}$$

in the right-hand side and the optimal value $f^*(\cdot)$ depends on the selection of \mathbf{p} . The equality and inequality constraints $\mathbf{x} \mapsto k(\mathbf{x})$ and $\mathbf{x} \mapsto g(\mathbf{x})$ cover the remaining constraints of (3.8), i.e., $X(0) = X_0$, (3.7), (CLI)–(EI), (2.2)–(2.4), $s(i), \mu(i) \in [0, 1]$ for $i \in \{0, \dots, N-1\}$ are encoded through appropriate equality and inequality constraints in the decision variable \mathbf{x} .

The Lagrangian corresponding to (3.9) with Lagrange multipliers $(\lambda_E, \lambda_C, \delta, \kappa)$ is defined as

$$\mathcal{L}(\mathbf{x}, \lambda_E, \lambda_C, \delta, \kappa; \mathbf{p}) = f(\mathbf{x}) - \left(\sum_{j=0}^{N-1} \lambda_{E_j} (p_{E_j} - h_{E_j}(\mathbf{x})) + \lambda_{C_j} (p_{C_j} - h_{C_j}(\mathbf{x})) \right) - \delta^T k(\mathbf{x}) - \kappa^T g(\mathbf{x}). \quad (3.10)$$

If \mathbf{x}^* and $(\lambda_E^*, \lambda_C^*, \delta^*, \kappa^*)$ are an optimal solution-Lagrange multiplier pair (see [27, Sec. 5.1.3], for example) then the optimal value function satisfies

$$f^*(\mathbf{p}) = \mathcal{L}(\mathbf{x}^*(\mathbf{p}), \lambda_E^*(\mathbf{p}), \lambda_C^*(\mathbf{p}), \delta^*(\mathbf{p}), \kappa^*(\mathbf{p}); \mathbf{p}) = f(\mathbf{x}^*(\mathbf{p})). \quad (3.11)$$

Moreover, under appropriate conditions (second order sufficiency conditions [27, Prop. 3.3.2], for example) the partial derivatives of the optimal value function in a neighborhood around $\mathbf{p} = 0$ satisfy

$$\begin{aligned} \frac{\partial f^*}{\partial p_{E_i}}(\mathbf{p}) &= \frac{\partial \mathcal{L}}{\partial p_{E_i}}(\mathbf{x}^*(\mathbf{p}), \lambda_E^*(\mathbf{p}), \lambda_C^*(\mathbf{p}), \delta^*(\mathbf{p}), \kappa^*(\mathbf{p}); \mathbf{p}) = -\lambda_{E_i}^*(\mathbf{p}), \\ \frac{\partial f^*}{\partial p_{C_i}}(\mathbf{p}) &= \frac{\partial \mathcal{L}}{\partial p_{C_i}}(\mathbf{x}^*(\mathbf{p}), \lambda_E^*(\mathbf{p}), \lambda_C^*(\mathbf{p}), \delta^*(\mathbf{p}), \kappa^*(\mathbf{p}); \mathbf{p}) = -\lambda_{C_i}^*(\mathbf{p}), \end{aligned} \quad (3.12)$$

for all $i \in \{0, \dots, N-1\}$, according to [27, Prop. 3.3.3].

While in the optimal value function necessarily the terms containing Lagrange multipliers vanish and only the objective function is important according to (3.11), in $\nabla_{\mathbf{p}} f^*(\mathbf{p})$ in (3.12) it turns out that the dependency of $(\mathbf{x}^*, \lambda_E^*, \lambda_C^*, \delta^*, \kappa^*)$ on \mathbf{p} can be ignored when computing the gradient through the Lagrangian (3.10). Thus, the Lagrange multipliers λ_E , λ_C are a measure of the sensitivity of the objective function with respect to changes in the right-hand side of the equality constraints. In economics, this sensitivity is also known under the name shadow price as outlined in [28, Chapter 3.3.4], [29, Chapter 3.5], for example.

These observations finally allow us to return to (3.3). If we recall the connection between (3.8) and (3.9) we observe that

$$\lambda_{E_j}^* = -\frac{\partial W^*}{\partial E^*(j)}, \quad \text{and} \quad \lambda_{C_j}^* = -\frac{\partial W^*}{\partial C^*(j)} \quad (3.13)$$

is satisfied for all $j \in \{0, \dots, N-1\}$. Thus according to (3.3) the SCC is given by

$$S_{CC}(j) = -1000 \frac{\lambda_{E_j}^*}{\lambda_{C_j}^*}. \quad (3.14)$$

Using (3.14), $S_{CC}(j)$ can be computed for all $j \in \{0, \dots, N-1\}$ by solving only a single optimization problem (3.8). In addition, unlike (3.5), Equation (3.14) is not an approximation of (3.3) but an exact expression. However, the pulse experiments might be more intuitive and from an interpretation point of view easier to understand than the connection to the Lagrange multipliers.

3.3. SCC calculation based on the discounted damages stream

In this section we present a third way to compute the SCC. This is based on the net present value, which, as described in the introduction, has been used in much of the climate economics research but not in a consistent manner. Here, we will show that a time-varying Ramsey discount factor, accounting for a time-varying growth rate of consumption, is required for consistency between calculating the SCC as a discounted damages stream and calculating the SCC as a ratio of marginal prices. The result relies on the assumption that (2.9) satisfies second order sufficiency conditions. The precise definition and a corresponding result are reported in Theorem 2 in Appendix A.

Theorem 1. Consider the optimal control problem (2.9) and assume that (2.9) satisfies second order sufficiency conditions. Then the SCC at time $j \in \{0, \dots, N-1\}$ is given by

$$S_{CC}(j) = -1000 \sum_{i=0}^{N-1} \frac{\partial C^*(i)}{\partial E^*(j)} \delta_{j,i}, \quad (3.15)$$

where the Ramsey discount factor between periods j and $i \in \{0, \dots, N-1\}$ is defined as

$$\delta_{j,i} := \frac{1}{(1+\rho)^{\Delta(i-j)}} \cdot \frac{1}{(1+g_{j,i})^\alpha}, \quad (3.16)$$

and the growth rate of per capita consumption up to period i is defined as

$$g_{j,i} := \frac{C^*(i)/L(i)}{C^*(j)/L(j)} - 1. \quad (3.17)$$

A proof of Theorem 1 is given in Appendix B. Based on Theorem 1 the SCC can again be computed using a finite difference approximation. While $\delta_{j,i}$ in (3.16) can be computed explicitly, the partial derivative $\frac{\partial C^*(i)}{\partial E^*(j)}$ in (3.15) needs to be approximated. We can thus perform two experiments to approximate the SCC at time $j \in \{0, \dots, N-1\}$ using Equation (3.15):

1. A baseline solution of (2.9) that generates a welfare value W^* corresponding to $E_{\text{Pulse}}(i) = 0$ and $C_{\text{Pulse}}(i) = 0$ for all $i = 0, \dots, N-1$ and additionally returns the nominal consumption pathway $C^*(i)$ for $i = 0, \dots, N-1$.
2. A second solution of (2.9) with a small non-zero perturbation on emissions at time $j \in \{0, \dots, N-1\}$, i.e., $E_{\text{Pulse}}(j) \neq 0$. This yields a second (optimal) welfare value $W^*[E_{\text{Pulse}}(j)]$ and additionally a second (optimal) consumption pathway $C^*[E_{\text{Pulse}}(j)](i)$ for $i = 0, \dots, N-1$.

With the resulting pathways, we can calculate the discount rate $\delta_{j,i}$ and we can approximate the partial derivatives, similar to (3.4), by $(C^*[E_{\text{Pulse}}(j)](i) - C^*(i))/(E_{\text{Pulse}}(j))$. The social cost of carbon can then be approximated by

$$S_{CC}(j) \approx -\frac{1000}{E_{\text{Pulse}}(j)} \sum_{i=0}^{N-1} (C^*[E_{\text{Pulse}}(j)](i) - C^*(i)) \delta_{j,i}. \quad (3.18)$$

The expression (3.18) has a familiar interpretation as the net present value (NPV) of the stream of damages due to an emissions pulse at time j , and is conventionally expressed in units of \$/tCO₂. Note, however, that in computing the NPV in (3.18), the discount factor $\delta_{j,i}$ is time-varying due to its dependence on the growth rate $g(\cdot)$ of per capita consumption; cf. (3.16). This marks a subtle, but potentially interesting point of departure between (3.18) and a large body of literature which applies the well-known, continuous-time Ramsey equation for the discount rate $r = \rho + \alpha \cdot g$ for constant g , to the discrete-time setting (see [30], for example).

Remark 2. Using the Taylor expansion

$$C^*[E_{\text{Pulse}}(j)](i) = C^*(i) + \frac{\partial C^*}{\partial E^*(j)} E_{\text{Pulse}}(j) + O(E_{\text{Pulse}}(j)^2), \quad (3.19)$$

i.e.,

$$\frac{\partial C^*}{\partial E^*(j)} = \frac{C^*[E_{\text{Pulse}}(j)](i) - C^*(i)}{E_{\text{Pulse}}(j)} + O(E_{\text{Pulse}}(j)),$$

it follows that (3.18) converges with order $O(E_{\text{Pulse}}(j))$ to $S_{\text{CC}}(j)$, similar to the first order convergence of (3.5).

4. Numerical Results

In this section we discuss three aspects in the calculation of the SCC using the DICE model. For the calculations we use the standard setting of the DICE-2016R model (see [24, Table 2] for the parameter selection) if not specified differently. For the numerical calculations we use the implementation [31]. While we use a horizon of $N = 100$, consistent with the parameters in DICE-2016R, we only focus on the first time steps in the visualizations.

Before we focus on the calculation of the SCC, we recall one of its illustrative definitions and visualizations used in the literature and highlight the anticipatory response of solutions of the optimal control problem (2.9) when emission or consumption impulses are added at different time steps. In [4] the SCC is illustrated as “the economic cost caused by an additional ton of carbon dioxide emissions” which is usually illustrated by adding a pulse to the first time step in the future [4, Figure 1]. In particular, with respect to [4, Figure 1], if time step 1 indicates the initial conditions, an additional ton of carbon dioxide emissions is added at time step 2. Under these assumptions, baseline emission and consumption pathways are compared with perturbed emission and consumption pathways. The baseline emissions and consumption pathways using the DICE-2016R model are visualized in Figure 1.




Figure 1. Optimal (baseline) emissions and consumption pathways using the DICE-2016R model.

By adding the additional tonne at the earliest possible time in the future, one might get the misleading impression that a perturbation $E_{\text{Pulse}}(j)$, $j \in \{0, \dots, N - 1\}$, only affects emissions and consumption at time steps $k \in \{j, \dots, N - 1\}$. This is however not the case as shown in Figure 2, where a pulse is added at time $j = 1$, $j = 10$ and $j = 20$, respectively. In Figure 2 the difference between the baseline so-




Figure 2. Impact of an additional tonne of emissions at different time steps with respect to the optimal emissions and consumption pathways.

lution, i.e., $E_{\text{Pulse}}(j) = C_{\text{Pulse}}(j) = 0$ for all $j \in \{0, \dots, N-1\}$, and perturbed solutions with $C_{\text{Pulse}}(j) = 1$ for $j = 1$, $j = 10$ and $j = 20$ are shown. A pulse at time $j \in \{0, \dots, N-1\}$ has an impact on the whole pathways $C[E_{\text{Pulse}}(j)](\cdot)$ and $E[E_{\text{Pulse}}(j)](\cdot)$ due to the optimization involved in their computation. This does also become clear from the definition of the partial derivative (3.2) used in the formula for the SCC: for computing this derivative, first the pulse is added and then the optimization is performed, implying that the optimizer is aware of the pulse at time j when performing the optimization for the times $i = 0, \dots, j-1$. In other words, if there is a change of emissions in the future, which importantly by assumption is known, then the change is anticipated and encoded in the optimal control problem (2.9). This explains the non-negligible difference in $C[C_{\text{Pulse}}(20)](i)$ and $C(i)$ and in $E[C_{\text{Pulse}}(i)](20)$ and $E(i)$, respectively, for $i < 20$, for example.

As a next step, we show that all three methods discussed in Section 3 essentially lead to the same SCC value. Figure 3 compares the three methods with each other, where for the pulse experiments the values $C_{\text{Pulse}}(j) = 1$ and $E_{\text{Pulse}}(j) = 1$, $j \in \{0, \dots, N-1\}$, were used for the finite difference approximations (which corresponds to a one tonne impulse in the implementation). On the scale on the left of Figure 3, the SCC values are indistinguishable. However, the difference between the Lagrange multiplier method and the finite difference approximation (3.5) and the formula (3.18) derived in Theorem 1 appears to grow nonlinearly with $j \in \{0, \dots, N-1\}$ as visualized in Figure 3 on the right.

In Figure 4 the error in the SCC calculation with respect to E_{Pulse} and C_{Pulse} are shown. In particular, the impulses $E_{\text{Pulse}}(j)$ and $C_{\text{Pulse}}(j)$ are varied from 0.01 to 5 for different values of $j \in \{0, \dots, N-1\}$. As expected from Remarks 1 and 2, the error grows linearly with E_{Pulse} and C_{Pulse} , respectively.

Remark 3. *While in theory the SCCs computed with the three methods coincide, the precise numerical values we obtain are sensitive with respect to E_{Pulse} and C_{Pulse} , the chosen optimization algorithm and the parameter selection of the optimization algorithm. Figure 5 shows the same results as Figure 4 with the only difference that the horizon $N = 100$ has been replaced with $N = 150$. In this case the parameter selection used in the IPOPT solver does not lead to reliable results for all impulse perturbations.*

As a final visualization we show the change of the SCC with respect to the discount rate in Figure




Figure 3. Comparison of the SCC calculation with three different methods discussed in Section 3.




Figure 4. Error in the calculation of the SCC. In general, the error changes linear with the pulse. However, some values of the perturbations lead to numerical problems.

6, i.e., $\frac{d}{d\rho} S_{CC}(j)$ is calculated and shown in Figure 6. Here $\frac{d}{d\rho} S_{CC}(j)$ is approximated through a finite difference approximation

$$\frac{d}{d\rho} S_{CC}(j) \approx \frac{1}{\Delta\rho} (S_{CC}[\rho + \Delta\rho](j) - S_{CC}[\rho](j)),$$

where $S_{CC}[\rho + \Delta\rho](j)$ and $S_{CC}[\rho](j)$ are calculated using the Lagrange multiplier method presented in Section 3.2.

5. Conclusions

In this paper, we have investigated and compared three different methods for the computation of the SCC in integrated assessment models. While our discussion is focused on the DICE model, it is equally relevant for other models including PAGE and FUND, to obtain a characterization of the SCC that is consistent among the literature. While the first two methods, which rely on perturbations of a baseline model and on dual variables of an appropriately defined optimal control problem, respectively, are commonly known and used, the third method relies on a novel formula stated in Theorem 1. The novel formula allows a direct analysis of the impact of the discount factor in the calculation of the NPV and the SCC through Equation (3.16), which subsequently provides insights in SCC values stemming from different integrated assessment models.




Figure 5. Error in the calculation of the SCC similar to Figure 4 with the only difference that $N = 100$ has been replaced by $N = 150$. Because of the increased horizon, the numerical results are not reliable.




Figure 6. Sensitivity of the SCC with respect to the discount rate.

Appendix

A. Optimality conditions for constrained and parametric optimization

In this section we recapitulate optimality conditions for constrained optimization problems from [27], which are needed for the proof of the main Theorem 1 in Appendix B. In particular, for the convenience of the reader, we rewrite results for minimization problems in terms of maximization problems discussed in this paper. We consider optimization problems of the form

$$\begin{aligned} f^* &= \max_x f(x) \\ &\text{subject to } h(x) = 0, \\ &\quad g(x) \leq 0. \end{aligned} \tag{A.1}$$

where $x \in \mathbb{R}^n$, $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $h : \mathbb{R}^n \rightarrow \mathbb{R}^m$, and $g : \mathbb{R}^n \rightarrow \mathbb{R}^p$ for $n, m, p \in \mathbb{N}$. With the Lagrange multipliers $\lambda \in \mathbb{R}^m$ and $\mu \in \mathbb{R}_{\geq 0}^p$, the Lagrangian corresponding to (A.1) is defined as

$$\mathcal{L}(x, \lambda, \mu) = f(x) - \lambda^T h(x) - \mu^T g(x).$$

Additionally, for $x \in \mathbb{R}^n$ we define

$$A(x) = \{i \in \{1, \dots, p\} \mid g_i(x) = 0\}$$

as the corresponding active set with respect to the inequality constraints g .

Theorem 2 (Second order Sufficiency Conditions, [27, Prop. 3.3.2]). *Consider the optimization problem (A.1) and assume that f , h and g are twice continuously differentiable. Let $x^* \in \mathbb{R}^n$, $\lambda^* \in \mathbb{R}^m$ and $\mu^* \in \mathbb{R}^m$ satisfy*

$$\begin{aligned} \nabla_x \mathcal{L}(x^*, \lambda^*, \mu^*) &= 0, & h(x^*) &= 0, & g(x^*) &\leq 0, \\ \mu_j^* &\geq 0, & \forall j &\in \{1, \dots, p\} \\ \mu_j^* &= 0, & \forall j &\in \{1, \dots, p\} \setminus A(x^*) \\ y^T \nabla_{xx}^2 \mathcal{L}(x^*, \lambda^*, \mu^*) y &< 0 \end{aligned}$$

for all $y \neq 0$ such that

$$\nabla h_i(x^*)^T y = 0, \quad \forall i = 1, \dots, m, \quad \nabla g_i(x^*)^T y = 0 \quad \forall i \in A(x^*).$$

Finally, assume that

$$\mu_j^* > 0 \quad \forall j \in A(x^*).$$

Then x^* is a strict local maximum of (A.1) and we say that (A.1) satisfies the second order sufficient condition at x^* .

Additionally we will need the following result for parameter dependent optimization problems where the parameter only appears in the objective function $f : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$.

Theorem 3 ([32, Corollary 3.4.2]). *Consider the parameter dependent optimization problem*

$$\begin{aligned} f^*(p) &= \max_x f(x, p) \\ &\text{subject to } h(x) = 0 \\ &\quad g(x) \leq 0 \end{aligned} \tag{A.2}$$

with Lagrangian

$$\mathcal{L}(x, \lambda, \mu, p) = f(x, p) - \lambda^T h(x) - \mu^T g(x),$$

Lagrange multipliers λ , μ and parameter $p \in \mathbb{R}$ in the objective function. Assume that the second order sufficiency conditions of Theorem 2 are satisfied. Then, in a neighborhood around $p = 0$ it holds that

$$f^*(p) = \mathcal{L}(x^*(p), \lambda^*(p), \mu^*(p), p), \tag{A.3}$$

$$\nabla_p f^*(p) = \nabla_p \mathcal{L}(x^*(p), \lambda^*(p), \mu^*(p), p), \tag{A.4}$$

$$\nabla_p^2 f^*(p) = \nabla_x^2 \mathcal{L}(x^*(p), \lambda^*(p), \mu^*(p), p) \nabla_p x^*(p) + \nabla_p^2 \mathcal{L}(x^*(p), \lambda^*(p), \mu^*(p), p). \tag{A.5}$$

B. Proof of Theorem 1

To prove Theorem 1 we will make use of results in [32] and the results summarized in Appendix A. Before we begin with the actual proof we note that for

$$\mathbf{x} = \begin{bmatrix} \mathbf{s} \\ \boldsymbol{\mu} \end{bmatrix}$$

the optimization problems (2.9) can be written in the general form

$$\begin{aligned} f^*(p) &= \max_{\mathbf{x}} f(\mathbf{x}, p) \\ &\text{subject to } g(\mathbf{x}) \leq 0, \end{aligned} \quad (\text{B.1})$$

where the equality constraints are eliminated and the box constraints are captured in the function g . Here, the parameter p represents the pulses $E_{\text{Pulse}}(j)$ and $C_{\text{Pulse}}(j)$, or equivalently, p represents a change in $C(j)$ and $E(j)$, for $j \in \{0, \dots, N-1\}$. Since we only consider the perturbation with respect to C or E and only with respect to one $j \in \{0, \dots, N-1\}$ at a time, we assume that p is one dimensional. With these definitions we can use Theorem 3 (without the function h) in the following. Additionally we note that according to the definition of the functions involved in the optimal control problem (2.9), the functions f and g are twice continuously differentiable.

Proof of Theorem 1: We begin the proof by rewriting the partial derivatives

$$\frac{\partial W^*}{\partial C^*(j)} \quad \text{and} \quad \frac{\partial W^*}{\partial E^*(j)}$$

for $j \in \{0, \dots, N-1\}$. To this end we define

$$c_i = c_i(C(i)) = 1000 \frac{C(i)}{L(i)}, \quad (\text{B.2})$$

which is a scaling of $C(i)$ through the known parameter $L(i)$, $i \in \{0, \dots, N-1\}$. Thus the utility U defined in (2.7) can be written as a function of c_i ,

$$U(c_i) = L(i) \frac{c_i^{1-\alpha} - 1}{1-\alpha}. \quad (\text{B.3})$$

For the derivative we obtain

$$\frac{\partial U}{\partial c_i}(c_i) = L(i) c_i^{-\alpha}, \quad (\text{B.4})$$

and with respect to the original variables it holds that

$$\frac{\partial U}{\partial C(i)}(c_i) = \frac{\partial U(c_i)}{\partial c_i} \frac{\partial c_i(C(i))}{\partial C(i)} = L(i) c_i^{-\alpha} \frac{1000}{L(i)} = 1000 c_i^{-\alpha}. \quad (\text{B.5})$$

With Theorem 3, and in particular with (A.4) for $p = E_{\text{Pulse}}(j)$, $j \in \{0, \dots, N-1\}$, it holds that

$$\frac{\partial W^*}{\partial E^*(j)} = \frac{\partial}{\partial E^*(j)} \left(\sum_{i=0}^{N-1} (1-\rho)^{-\Delta i} U(c_i^*) \right).$$

Using the notation

$$\gamma_i = (1 - \rho)^{-\Delta i}$$

to shorten the expressions and the equation derived in Equation (B.5) the right-hand side can be written in the form

$$\frac{\partial W^*}{\partial E^*(j)} = \sum_{i=0}^{N-1} \gamma_i \frac{\partial U}{\partial E^*(j)}(c_i^*) = \sum_{i=0}^{N-1} \gamma_i \frac{\partial U(c_i^*)}{\partial C(i)} \frac{\partial C^*(i)}{\partial E(j)} = \sum_{i=0}^{N-1} \gamma_i 1000(c_i^*)^{-\alpha} \frac{\partial C^*(i)}{\partial E(j)}. \quad (\text{B.6})$$

Similarly, the application of Equation (A.3) in Theorem 3, where p represents a change in $C(j)$, $j \in \{0, \dots, N-1\}$, leads to the expression

$$\begin{aligned} \frac{\partial W^*}{\partial C^*(j)} &= \frac{\partial}{\partial C^*(j)} \left(\sum_{i=0}^{N-1} \gamma_i U(c_i^*) \right) = \sum_{i=0}^{N-1} \gamma_i \frac{\partial U(c_i^*)}{\partial C^*(j)} = \sum_{i=0}^{N-1} \gamma_i \frac{\partial U(c_i^*)}{\partial C^*(i)} \frac{\partial C^*(i)}{\partial C^*(j)} \\ &= \sum_{i=0}^{N-1} \gamma_i 1000(c_i^*)^{-\alpha} \frac{\partial C^*(i)}{\partial C^*(j)}. \end{aligned} \quad (\text{B.7})$$

As a next step, we derive Taylor expansions of the optimal value function W^* . In particular for parameters p_i representing a change in $C^*(j)$, (i.e., a pulse $C_{\text{Pulse}}(i)$) for $i \in \{0, \dots, N-1\}$, we derive Taylor expansions in a neighborhood of $p_i = 0$. Using the definition of c_i in Equation (B.2), the consumption (2.6) can be equivalently written as

$$c_i = \frac{10^3}{L(i)} \left(\frac{1}{1 + a T_{\text{AT}}(i)^2} \right) \left(1 - \theta_1(i) \mu(i)^{\theta_2} \right) A(i) K(i)^\gamma L(i)^{1-\gamma} (1 - s(i)) + \frac{10^3}{L(i)} C_{\text{Pulse}}(i). \quad (\text{B.8})$$

Moreover, with the definition $c_{\text{Pulse}}(i) = \frac{10^3}{L(i)} C_{\text{Pulse}}(i)$ and $c_i = \tilde{c}_i + c_{\text{Pulse}}(i)$, Equation (B.8) can be written as

$$\tilde{c}_i = \frac{L(i)}{10^3} \left(\frac{1}{1 + a T_{\text{AT}}(i)^2} \right) \left(1 - \theta_1(i) \mu(i)^{\theta_2} \right) A(i) K(i)^\gamma L(i)^{1-\gamma} (1 - s(i)) \quad (\text{B.9})$$

and the utility function can be written as

$$U[c_{\text{Pulse}}(i)](\tilde{c}_i) = L(i) \frac{(\tilde{c}_i + c_{\text{Pulse}}(i))^{1-\alpha} - 1}{1 - \alpha}. \quad (\text{B.10})$$

Thus, instead of the parameter $C_{\text{Pulse}}(j)$ in the consumption (2.6) we can use a scaled parameter directly in the utility function (2.7) and the optimal value function satisfies

$$W^*[p_i] = \gamma_i U(\tilde{c}_i^*[p_i] + p_i) + \sum_{\substack{k=0 \\ k \neq i}}^N \gamma_k U(\tilde{c}_k^*[p_i])$$

with $p_i = c_{\text{Pulse}}(i)$. According to Theorem 3 (using Equation (A.5)) the optimal value function is twice continuously differentiable with respect to the parameter p_i , and using Theorem 3, Equation (A.4), the Taylor approximation is given by

$$W^*[p_i] = W^*[0] + \gamma_i L(i) (\tilde{c}_i^*[0])^{-\alpha} p_i + R_i(p_i)$$

where

$$R_i(p_i) = \frac{1}{2} \frac{d^2}{dp_i^2} W^*[\xi] p_i^2 \quad (\text{B.11})$$

for $\xi \in [0, p_i]$. For $i, j \in \{0, \dots, N-1\}$ we take specific perturbations $p_i \neq 0, p_j \neq 0$ with the property $W^*[p_i] = W^*[p_j]$ so that the Taylor approximation satisfies

$$W^*[0] + \gamma_i L(i)(\tilde{c}_i^*[0])^{-\alpha} p_i + R_i(p_i) = W^*[0] + \gamma_j L(j)(\tilde{c}_j^*[0])^{-\alpha} p_j + R_j(p_j). \quad (\text{B.12})$$

A p_j satisfying (B.12) can be found for each sufficiently small p_i because the terms in front of p_i and p_j in (B.12) are not zero due to the assumption that (2.9) satisfies the second order sufficiency conditions. In fact, $\gamma_j L(j)(\tilde{c}_j^*[0])^{-\alpha} = 0$ implies that $s^*(i) = 1$ or $\theta_1(i)\mu^*(i)^{\theta_2} = 1$ (see Equation (B.9)). If $s^*(i) = 1$ then $\mu^*(i) \in [0, 1]$ can be selected arbitrarily and if $\theta_1(i)\mu^*(i)^{\theta_2} = 1$ then $s^*(j) \in [0, 1]$ can be selected arbitrarily, i.e., the optimal solution of (2.9) is not characterized through a strict local maximum, which contradicts Theorem 2. Under these assumptions stated above, p_j is a function of p_i and $\lim_{p_i \rightarrow 0} p_j(p_i) = 0$ (since W^* is continuously differentiable). Rearranging terms yields

$$\frac{p_j}{p_i} = \frac{\gamma_i L(i)(\tilde{c}_i^*[0])^{-\alpha}}{\gamma_j L(j)(\tilde{c}_j^*[0])^{-\alpha}} + \frac{R_i(p_i) - R_j(p_j)}{p_i \gamma_j L(j)(\tilde{c}_j^*[0])^{-\alpha}} \quad (\text{B.13})$$

and allows us to investigate the limit $p_i \rightarrow 0, p_i \neq 0$, (i.e., $p_j(p_i) \rightarrow 0$). We observe that

$$\lim_{p_i \rightarrow 0} \frac{p_j(p_i)}{p_i} = \lim_{p_i \rightarrow 0} \frac{p_j(p_i) - p_j(0)}{p_i - 0} = \frac{dp_j}{dp_i}(0),$$

and since the remainder in Equation (B.11) satisfies $R_i(p_i) = O(p_i^2)$ we obtain

$$\lim_{p_i \rightarrow 0} \frac{1}{p_i} R_i(p_i) = 0.$$

To be able to neglect the second term in the right-hand side of Equation (B.13), it is left to show that $\frac{1}{p_i} R_j(p_j(p_i))$ converges to zero for $p_i \rightarrow 0$. With the definition of the remainder $R_j(p_j) = \frac{1}{2} \frac{d^2}{dp_j^2} W^*[\xi] p_j^2$ in Equation (B.11) the limit satisfies

$$\begin{aligned} \lim_{p_i \rightarrow 0} \left(\frac{1}{p_i} R_j(p_j(p_i)) \right) &= \lim_{p_i \rightarrow 0} \left(\frac{1}{2} \frac{d^2}{dp_j^2} W^*[\xi(p_i)] \frac{p_j(p_i)}{p_i} p_j(p_i) \right) \\ &= \frac{1}{2} \frac{d^2}{dp_j^2} W^*[\xi(0)] \frac{dp_j}{dp_i}(0) p_j(0) \\ &= \frac{1}{2} \frac{d^2}{dp_j^2} W^* \frac{dp_j}{dp_i}(0) \cdot 0 = 0. \end{aligned}$$

Consequently, using Equation (B.4) together with $\tilde{c}_i = c_i$ in the case $p_i = 0$, it holds that

$$\lim_{p_i \rightarrow 0} \frac{p_j}{p_i} = \gamma_{i-j} \frac{L(i)(c_i^*)^{-\alpha}}{L(j)(c_j^*)^{-\alpha}} = \gamma_{i-j} \frac{\frac{\partial U}{\partial c_i}(c_i^*)}{\frac{\partial U}{\partial c_j}(c_j^*)}. \quad (\text{B.14})$$

To return to the original variables $C(i)$, we recall the coordinate transformation (B.2), and use the notation $P_i = C_{\text{Pulse}}(j) = \frac{L(i)}{10^3} p_i$.

With the above computations we are finally in the position to derive the partial derivative of $C^*(i)$ with respect to $C^*(j)$ for $i, j \in \{0, \dots, N-1\}$. In particular, using the chain rule and the definition of the derivative, it holds that

$$\frac{\partial C^*(i)}{\partial C^*(j)} = \frac{\frac{\partial W^*}{\partial C^*(j)}}{\frac{\partial W^*}{\partial C^*(i)}} = \frac{\lim_{P_j \rightarrow 0} \frac{1}{P_j} (W^*[P_j] - W^*)}{\lim_{P_i \rightarrow 0} \frac{1}{P_i} (W^*[P_i] - W^*)}.$$

Using the relation between p_i and P_i , this expression can be equivalently written as

$$\frac{\partial C^*(i)}{\partial C^*(j)} = \frac{\lim_{p_j \rightarrow 0} \frac{10^3}{L(j)p_j} (W^*[p_j] - W^*)}{\lim_{p_i \rightarrow 0} \frac{10^3}{L(i)p_i} (W^*[p_i] - W^*)} = \lim_{p_i \rightarrow 0} \frac{L(i)p_i}{L(j)p_j} \frac{(W^*[p_j] - W^*)}{(W^*[p_i] - W^*)}.$$

Finally, with the result derived in Equation (B.14), it holds that

$$\frac{\partial C^*(i)}{\partial C^*(j)} = \lim_{p_i \rightarrow 0} \frac{L(i)p_i}{L(j)p_j(p_i)} = \gamma_{j-i} \frac{(c_j^*)^{-\alpha}}{(c_i^*)^{-\alpha}} \quad (\text{B.15})$$

where the second term is equal to one since $W^*[p_i] = W^*[p_j(p_i)]$ according the assumption made at the beginning of the derivation of (B.14) in (B.12).

To complete the proof we combine the results so far in the definition of $S_{\text{CC}}(j)$. With Equations (B.6) and (B.7) the SCC can be expressed as

$$S_{\text{CC}}(j) = -1000 \frac{\partial W^*/\partial E(j)}{\partial W^*/\partial C(j)} = -1000 \frac{\sum_{i=0}^{N-1} \gamma_i 1000 (c_i^*)^{-\alpha} \frac{\partial C^*(i)}{\partial E^*(j)}}{\sum_{i=0}^{N-1} \gamma_i 1000 (c_i^*)^{-\alpha} \frac{\partial C^*(i)}{\partial C^*(j)}}$$

and which becomes

$$S_{\text{CC}}(j) = -1000 \frac{\sum_{i=0}^{N-1} \gamma_i (c_i^*)^{-\alpha} \frac{\partial C^*(i)}{\partial E^*(j)}}{\sum_{i=0}^{N-1} \gamma_i (c_i^*)^{-\alpha} \gamma_{j-i} \frac{(c_j^*)^{-\alpha}}{(c_i^*)^{-\alpha}}}$$

through the identity (B.15). Finally, reformulating individual terms and recalling the definition of δ_{ji} in Theorem 1 completes the proof:

$$S_{\text{CC}}(j) = -1000 \frac{\sum_{i=0}^{N-1} \gamma_i (c_i^*)^{-\alpha} \frac{\partial C^*(i)}{\partial E^*(j)}}{\sum_{i=0}^{N-1} \gamma_j (c_j^*)^{-\alpha}} = -1000 \sum_{i=0}^{N-1} \gamma_{i-j} \frac{(c_i^*)^{-\alpha}}{(c_j^*)^{-\alpha}} \frac{\partial C^*(i)}{\partial E^*(j)} = -1000 \sum_{i=0}^{N-1} \frac{\partial C^*(i)}{\partial E^*(j)} \delta_{ji}.$$

□

C. Order of Convergence

As a last result in this paper we show that (3.5) converges with order $O(C_{\text{Pulse}}(j) + E_{\text{Pulse}}(j))$ to $S_{\text{CC}}(j)$. Using the notation introduced in (3.4) and (3.5) the SCC satisfies

$$S_{\text{CC}}(j) = -1000 \frac{\frac{\partial W^*}{\partial E^*(j)}}{\frac{\partial W^*}{\partial C^*(j)}} = -1000 \frac{\frac{W^*[E_{\text{Pulse}}(j)] - W^*}{E_{\text{Pulse}}(j)} + \Phi_E(E_{\text{Pulse}}(j))}{\frac{W^*[C_{\text{Pulse}}(j)] - W^*}{C_{\text{Pulse}}(j)} + \Phi_C(C_{\text{Pulse}}(j))} \quad (\text{C.1})$$

$$= -1000 \frac{C_{\text{Pulse}}(j)}{E_{\text{Pulse}}(j)} \cdot \frac{W^*[E_{\text{Pulse}}(j)] - W^*}{W^*[C_{\text{Pulse}}(j)] - W^*} - 1000 \Phi(E_{\text{Pulse}}(j), C_{\text{Pulse}}(j)). \quad (\text{C.2})$$

for unknown functions Φ_E , Φ_C and Φ . However, according to the Taylor expansions in (3.6) it holds that $\Phi_E(E_{\text{Pulse}}(j)) = O(E_{\text{Pulse}}(j))$ and $\Phi_C(C_{\text{Pulse}}(j)) = O(C_{\text{Pulse}}(j))$.

We investigate properties of the function Φ . To simplify the notation, we drop the index in the derivations, i.e., we investigate the equation

$$\frac{\frac{W^*[E]-W^*}{E} + \Phi_E(E)}{\frac{W^*[C]-W^*}{C} + \Phi_C(C)} = \frac{C}{E} \cdot \frac{W^*[E] - W^*}{W^*[C] - W^*} + \Phi(E, C),$$

or equivalently

$$\frac{C(W^*[E] - W^*) + E\Phi_E(E)}{E(W^*[C] - W^*) + C\Phi_C(C)} = \frac{C(W^*[E] - W^*) + \frac{E}{C}(W^*[C] - W^*)\Phi(E, C)}{E(W^*[C] - W^*)}.$$

We further simplify the notation and define $\Delta W_E^* = W^*[E] - W^*$ and $\Delta W_C^* = W^*[C] - W^*$. Then, multiplying by the denominator leads to the equation

$$\begin{aligned} \Delta W_C^* \Delta W_E^* + \Delta W_C^* E \Phi_E(E) &= (\Delta W_C^* + C\Phi_C(C)) \Delta W_E^* + \frac{E}{C} (\Delta W_C^* + C\Phi_C(C)) \Delta W_C^* \Phi(E, C) \\ &= \Delta W_C^* \Delta W_E^* + C\Phi_C(C) \Delta W_E^* + \frac{E}{C} (\Delta W_C^* + C\Phi_C(C)) \Delta W_C^* \Phi(E, C). \end{aligned}$$

After canceling and rearranging terms, we are left with the expression:

$$\Delta W_C^* E \Phi_E(E) - C\Phi_C(C) \Delta W_E^* = \frac{E}{C} (\Delta W_C^* + C\Phi_C(C)) \Delta W_C^* \Phi(E, C).$$

Since W^* is differentiable, i.e., the limits

$$\frac{\partial W^*}{\partial E^*(j)} = \lim_{\substack{E_{\text{Pulse}}(j) \rightarrow 0 \\ E_{\text{Pulse}}(j) \neq 0}} \frac{W^*[E_{\text{Pulse}}(j)] - W^*}{E_{\text{Pulse}}(j)}, \quad \frac{\partial W^*}{\partial C^*(j)} = \lim_{\substack{C_{\text{Pulse}}(j) \rightarrow 0 \\ C_{\text{Pulse}}(j) \neq 0}} \frac{W^*[C_{\text{Pulse}}(j)] - W^*}{C_{\text{Pulse}}(j)} \quad (\text{C.3})$$

exist, it holds that $\Delta W_E^* = O(E_{\text{Pulse}}(j))$ and $\Delta W_C^* = O(C_{\text{Pulse}}(j))$. Using these properties in the last equation simplifies the expression to

$$O(C)EO(E) + CO(C)O(E) = \frac{E}{C}(O(C) + CO(C))O(C)\Phi(E, C)$$

or equivalently

$$O(CE^2) + O(C^2E) = (O(CE) + O(C^2E))\Phi(E, C) = O(CE)\Phi(E, C).$$

In particular, Φ satisfies $\Phi(E, C) = O(C + E)$ proving the assertion.

References

1. D. Archer, M. Eby, V. Brovkin, A. Ridgwell, L. Cao, U. Mikolajewicz, K. Caldeira, K. Matsumoto, G. Munhoven, A. Montenegro, and K. Tokos. Atmospheric lifetime of fossil fuel carbon dioxide. *Annu. Rev. Earth Planet. Sci.*, 37:117–134, 2009.

2. M. Eby, K. Zickfeld, A. Montenegro, D. Archer, K.J. Meissner, and A.J. Weaver. Lifetime of anthropogenic climate change: Millennial time scales of potential CO₂ and surface temperature perturbations. *J. Clim.*, 22:2501–2511, 15 May 2009.
3. A. L. Marten, E. A. Kopits, C. W. Griffiths, S. C. Newbold, and A. Wolverton. Incremental CH₄ and N₂O mitigation benefits consistent with the US Government’s SC-CO₂ estimates. *Clim. Policy*, 15(2):272–298, 2015.
4. W. D. Nordhaus. Estimates of the social cost of carbon: Concepts and results from the DICE-2013R model and alternative approaches. *J. Assoc. Environ. Resour. Econ.*, 1(1/2):273–312, March 2014.
5. C. Hope and D. Newbery. Calculating the social cost of carbon. In M. Grubb, T. Jamasb, and M. G. Pollitt, editors, *Delivering a Low Carbon Electricity System: Technologies, Economics and Policy*, chapter 2, pages 31–63. Cambridge University Press, Cambridge, 2008.
6. D. Anthoff, R. S. J. Tol, and G. W. Yohe. Discounting for climate change. *Economics: the Open-Access, Open-Assessment e-Journal*, 3(2009-24):1–24, 9 June 2009.
7. W. D. Nordhaus and P. Sztorc. *DICE 2013R: Introduction and User’s Manual*, second edition, 31 October 2013.
8. F. P. Ramsey. A mathematical theory of saving. *Econ. J.*, 38(152):543–559, December 1928.
9. D. Anthoff and R.S.J. Tol. The uncertainty about the social cost of carbon: A decomposition analysis using FUND. *Clim. Change*, 117(3):515–530, April 2013.
10. D. Anthoff, R. S. J. Tol, and G. W. Yohe. Risk aversion, time preference, and the social cost of carbon. *Environ. Res. Lett.*, 4:1–7, 2009.
11. C. Hope. Critical issues for the calculation of the social cost of CO₂: why the estimates from PAGE09 are higher than those from PAGE2002. *Clim. Change*, 117:531–543, 2013.
12. T. Kögel. The rate of change of the social cost of carbon and the social planner’s Hotelling rule. *Economics: the Open-Access, Open-Assessment e-Journal*, pages 1–26, 14 August 2012. 2012-37.
13. A. L. Marten. Transient temperature response modeling in IAMs: The effects of over simplification on the SCC. *Economics: the Open-Access, Open-Assessment e-Journal*, pages 1–44, October 2011.
14. A. L. Marten and S. C. Newbold. Estimating the social cost of non-CO₂ GHG emissions: Methane and nitrous oxide. *Energy Policy*, 51:957–972, December 2012.
15. E. J. Moyer, M. D. Woolley, N. J. Matteson, M. J. Glotter, and D. A. Weisbach. Climate impacts on economic growth as drivers of uncertainty in the social cost of carbon. *J. Leg. Stud.*, 43(2):401–425, June 2014.
16. R. S. J. Tol. An analysis of mitigation as a response to climate change. Technical report, Copenhagen Consensus Center, 2009.
17. R. S. J. Tol. *Climate Economics: Economic Analysis of Climate, Climate Change, and Climate Policy*. Edward Elgar, 2014.
18. G. Economides, A. Papandreou, E. Sartzetakis, and A. Xepapadeas. *The Economics of Climate Change*. Bank of Greece, 2018.
19. S. C. Newbold, C. Griffiths, C. Moore, A. Wolverton, and E. Kopits. A rapid assessment model for understanding the social cost of carbon. *Clim. Change Econ.*, 4(1):1–40, 2013.

-
20. F. van der Ploeg. Discounting and climate policy, 2020. CESifo Working Paper no. 8441, Available at SSRN: <https://ssrn.com/abstract=3657977>.
 21. K. Arrow, M. Cropper, C. Gollier, B. Groom, G. Heal, R. Newell, W. Nordhaus, W. Pindyck, W. Pizer, P. Portney, T. Sterner, R. S. J. Tol, and M. Weitzman. Determining benefits and costs for future generations. *Science*, 341:349–350, 26 July 2013.
 22. B. Groom, C. Hepburn, P. Koundouri, and D. Pearce. Declining discount rates: The long and the short of it. *Environ. Resour. Econ.*, 32(4):445–493, December 2005.
 23. D. A. Farber. Gambling over growth: Economic uncertainty, discounting, and regulatory policy. *J. Leg. Stud.*, 44(2):S509–S528, June 2015.
 24. C. M. Kellett, S. R. Weller, T. Faulwasser, L. Grüne, and W. Semmler. Feedback and optimal control in climate economics. *Annu. Rev. Control*, 47:7–20, 2019.
 25. A. Wächter and L. T. Biegler. On the implementation of an interior-point filter line-search algorithm for large-scale nonlinear programming. *Mathematical programming*, 106:25–57, 2006.
 26. A. S. Drud. CONOPT—a large-scale GRG code. *ORSA Journal on computing*, 6(2):207–216, 1994.
 27. D. P. Bertsekas. *Nonlinear Programming*. Athena Scientific, 1999.
 28. D. Grass, J. P. Caulkins, G. Feichtinger, G. Tragler, and D. A. Behrens. *Optimal Control of Nonlinear Processes: With Applications in Drugs, Corruption, and Terror*. Springer, 2008.
 29. A. Seierstad and K. Sydsaeter. *Optimal Control Theory with Economic Applications*. Elsevier North-Holland, 1986.
 30. J. Y. Campbell and L. M. Viceira. *Strategic Asset Allocation: Portfolio Choice for Long-Term Investors*. Oxford University Press, 2002.
 31. T. Faulwasser, C. M. Kellett, and S. R. Weller. MPC-DICE: An open-source Matlab implementation of receding horizon solutions to DICE. In *Proc. 1st IFAC Workshop on Integrated Assessment Modeling for Environmental Systems*, Brescia, Italy, May 2018.
 32. A. V. Fiacco. *Introduction to Sensitivity and Stability Analysis in Nonlinear Programming*. Academic Press, 1983.