

Vector space partitions of $\text{GF}(2)^8$

Sascha Kurz

University of Bayreuth, 95440 Bayreuth, Germany
sascha.kurz@uni-bayreuth.de

February 2022

Abstract

A vector space partition \mathcal{P} of the projective space $\text{PG}(v-1, q)$ is a set of subspaces in $\text{PG}(v-1, q)$ which partitions the set of points. We say that a vector space partition \mathcal{P} has type $(v-1)^{m_{v-1}} \dots 2^{m_2} 1^{m_1}$ if precisely m_i of its elements have dimension i , where $1 \leq i \leq v-1$. Here we determine all possible types of vector space partitions in $\text{PG}(7, 2)$.

Keywords: Finite geometry, vector space partitions, divisible codes, linear codes

1 Introduction

We call the i -dimensional subspaces of the projective space $\text{PG}(v-1, q)$ i -spaces using the geometric names points, lines, planes, solids, and hyperplanes for 1-, 2-, 3-, 4-, and $(v-1)$ -spaces, respectively. A *vector space partition* \mathcal{P} of $\text{PG}(v-1, q)$ is a set of subspaces in $\text{PG}(v-1, q)$ which partitions the set of points. For a survey on known results we refer to [Hed12]. We say that a vector space partition \mathcal{P} has type $(v-1)^{m_{v-1}} \dots 2^{m_2} 1^{m_1}$ if precisely m_i of its elements have dimension i , where $1 \leq i \leq v-1$. The classification of the possible types of a vector space partition, given the parameters v and q , is an important and difficult problem. Based on [Hed86], the classification for the binary case $q = 2$ was completed for $v \leq 7$ in [EZSS⁺09]. Under the assumption $m_1 = 0$ the case $(q, v) = (2, 8)$ has been treated in [EZHS⁺10]. Here we complete the classification of the possible types of vector space partitions in $\text{PG}(7, 2)$. We will also briefly discuss the feasible types of vector space partitions in $\text{PG}(v-1, q)$ for all field sizes q and all dimensions $v \leq 5$.

Setting $[k]_q := \frac{q^k - 1}{q - 1}$ we can state that every k -space in $\text{PG}(v-1, q)$ consists of $[k]_q$ points. So, counting points gives the *packing condition*

$$\sum_{i=1}^{v-1} m_i \cdot [i]_q = [v]_q. \quad (1)$$

Another well-known condition uses the fact that an a -space and a disjoint b -space span an $(a+b)$ -space. So, we have

$$m_i \cdot m_j = 0 \quad (2)$$

for all $1 \leq i < j \leq v-1$ with $i+j > j$ and $m_i \leq 1$ for all $i > v/2$. These two conditions are sufficient to characterize all feasible types of vector space partitions in $\text{PG}(v-1, q)$ for $v \leq 4$.

Another condition stems from the fact that for an index $2 \leq j \leq v-1$ with $m_j > 0$ the set of points contained in the subspaces of dimension strictly less than j corresponds to a q^{j-1} -divisible linear code over \mathbb{F}_q of length

$$n = \sum_{i=1}^{j-1} m_i \cdot [i]_q, \quad (3)$$

see e.g. [KK20], where a linear code is called Δ -divisible if all of its codewords have a weight that is divisible by Δ . Non-existence results for q^r -divisible projective codes are e.g. discussed in [HKK18]. A recent survey can be found in [Kur21]. Some of these conditions are already contained in [Hed09] and used under the name *tail condition* in the literature on vector space partitions.

A few more necessary conditions for the existence of vector space partitions are stated in [LH12].

The notion of a vector space partition can be generalized in several directions. A λ -fold vector space partition of $\text{PG}(v-1, q)$ is a (multi-) set of subspaces such that every point P is covered exactly λ times, see e.g. [EZSS⁺11]. Another variant considers sets of subspaces such that every t -space is covered exactly once, see [HHKK19]. Vector space partitions of affine spaces have been considered in [BFIK22].

The remaining part of the paper is structured as follows. In Section 2 we introduce the necessary preliminaries. We deduce our main result – the classification of all possible types of vector space partitions of $\text{PG}(7, 2)$ – in Section 3. While several of the presented non-existence results for vector space partitions have purely theoretical proofs, others rely on extensive computer computations. It would be nice to see some of these calculations be replaced by pen-and-paper proofs. To top the paper off, we discuss the possible types of vector space partitions in $\text{PG}(v-1, q)$ for arbitrary field sizes q and dimensions $v \leq 5$ of the ambient space in Section 4.

2 Preliminaries

A vector space partition \mathcal{P} of $\text{PG}(v-1, q)$ is called *reducible* if there exists a proper subset \mathcal{Q} of the elements of \mathcal{P} whose points partition a proper subspace of $\text{PG}(v-1, q)$. If \mathcal{P} is not reducible we also speak of an irreducible vector space partition. We can easily construct reducible vector space partitions by starting from an arbitrary vector space partition and replacing an element with dimension at least 2 by its contained points.

A *multiset of points* in $\text{PG}(v-1, q)$ is a mapping χ from the set of points in $\text{PG}(v-1, q)$ to \mathbb{N} , so that $\chi(P)$ is the multiplicity of the point P . If $\chi(P) \leq 1$ for all points P , then we also speak of a set (of points) χ in $\text{PG}(v-1, q)$. The *support* of a multiset of points is the set of all points with non-zero multiplicity. We say that χ is Δ -divisible if $\sum_{P \not\leq H} \chi(P) \equiv 0 \pmod{\Delta}$ for each hyperplane H of $\text{PG}(v-1, q)$ and points P . In other words, the Hamming weights of the codewords of the \mathbb{F}_q -linear code C_χ associated with χ are divisible by Δ . By $\#\mathcal{M}$ we denote the cardinality of \mathcal{M} , i.e., the sum $\sum_P \mathcal{M}(P)$ of the multiplicities of all points P . If S is an arbitrary subspace, then by $\mathcal{M}(S)$ we denote the sum $\sum_{P \leq S} \mathcal{M}(P)$ of the point multiplicities of the points contained in S . For each $U \in \text{PG}(v-1, q)$ we denote by χ_U its characteristic function, i.e., $\chi_U(P) = 1$ iff $P \leq U$ and $\chi_U(P) = 0$ otherwise. It is an easy exercise to show that χ_U is $q^{\dim(U)-1}$ -divisible, which extends to multisets of subspaces:

Lemma 1. ([KK20, Lemma 11]) *Let \mathcal{U} be a multiset of subspaces in $\text{PG}(v-1, q)$ with dimension at least k . Then $\chi_{\mathcal{U}} := \sum_{U \in \mathcal{U}} \chi_U$ is q^{k-1} -divisible.*

If χ is Δ -divisible and $\chi(P) \leq \lambda$ for some constant $\lambda \in \mathbb{N}$ and all points P , then the λ -complement $\bar{\chi}$, defined by $\bar{\chi}(P) = \lambda - \chi(P)$ for all points P , is also Δ -divisible.

Corollary 2. *Let \mathcal{P} be a vector space partition of $\text{PG}(v-1, q)$ of type $(v-1)^{m_{v-1}} \dots 2^{m_2} 1^{m_1}$, then the set of points \mathcal{H}_k such that the corresponding element $A \in \mathcal{P}$, that contains the point, has dimension at most k is q^k -divisible if \mathcal{P} contains an element with dimension strictly larger than k .*

An exemplary implication is that no vector space partition of type $3^8 2^2 1^1$ in $\text{PG}(5, 2)$ exists since there is no 2-divisible multiset of points of cardinality 1, i.e., no 2-divisible binary code of effective length 1. In our setting of vector space partitions the maximum possible point multiplicity is 1, so that the corresponding codes are projective, i.e., generator matrices do not contain repeated columns. The possible effective lengths of projective binary Δ -divisible linear codes have been completely characterized for all $\Delta \in \{2, 4, 8\}$, see e.g. [HKK18, HKKW19] for proofs and further references:

Proposition 3. Let $n \in \mathbb{N}_{>0}$ be the effective length of a Δ -divisible binary projective linear code.

- (a) If $\Delta = 2$, then $n \geq 3$.
- (b) If $\Delta = 4$, then $n \in \{7, 8\}$ or $n \geq 14$.
- (c) If $\Delta = 8$, then $n \in \{15, 16, 30, 31, 32, 45, 46, 47, 48, 49, 50, 51\}$ or $n \geq 60$.

All those effective lengths can indeed be attained.

Definition 4. We say that a vector space partition \mathcal{P} has an s -supertail of type $a_1^{m_1} a_2^{m_2} \dots a_s^{m_s}$, where $a_1 > a_2 > \dots a_s \geq 1$, if \mathcal{P} contains exactly m_j elements of dimension a_j for all $1 \leq j \leq s$, there exists at least one element $A \in \mathcal{P}$ with $\dim(A) > a_1$, and for all elements $B \in \mathcal{P}$ with $\dim(B) \leq a_1$ there exists an index $1 \leq j \leq s$ with $\dim(B) = a_j$.

For the ease of notation we also allow the choice of $m_j = 0$ and just speak of a supertail of a certain type. From Proposition 3 we can directly conclude that certain types of supertails are impossible:

Corollary 5. Let \mathcal{P} be a vector space partition of $\text{PG}(v-1, 2)$, then \mathcal{P} cannot have a supertail of one of the following types:

- $1^1, 1^2$;
- $2^0 1^i$ for $i \in \{1, \dots, 6, 9, \dots, 13\}$, $2^1 1^i$ for $i \in \{0, \dots, 3, 6, \dots, 10\}$, $2^2 1^i$ for $i \in \{0, 3, \dots, 7\}$, $2^3 1^i$ for $i \in \{0, \dots, 4\}$, $2^4 1^i$ for $i \in \{0, 1\}$; and
- $3^a 2^b 1^c$ where $7a + 3b + c < 60$ and $7a + 3b + c \notin \{0, 15, 16, 30, 31, 32, 45, 46, 47, 48, 49, 50, 51\}$.

For literature on the supertail we refer e.g. to [HLNS13, NS18].

For small n the projective Δ -divisible \mathbb{F}_q -linear codes of effective length n can be exhaustively generated with software packages like e.g. **Q-Extension** [Bou07] or **LinCode** [BBK21]. Having the point sets at hand, we can check whether they can be partitioned into a certain number of planes, lines, and points, which excludes a few further supertail types. E.g. one can easily show that each q^2 -divisible multiset of cardinality $q^2 + q + 1$ over \mathbb{F}_q has to be the characteristic function of a plane, so that there is no supertail of type $2^2 1^{q^2 - q - 1}$ over \mathbb{F}_q . For enumeration results of projective binary Δ -divisible codes for $\Delta \in \{2, 4, 8\}$ we refer to [HHK⁺17]. Since the corresponding codes are computationally shown to be unique we have:

Lemma 6. Let \mathcal{S} be a Δ -divisible set of cardinality n over \mathbb{F}_2 .

- (a) If $(\Delta, n) = (2, 3)$, then \mathcal{S} is the characteristic function of a line.
- (b) If $(\Delta, n) = (4, 7)$, then \mathcal{S} is the characteristic function of a plane.
- (c) If $(\Delta, n) = (4, 14)$, then \mathcal{S} is the sum of the characteristic functions of two disjoint planes.
- (d) If $(\Delta, n) = (8, 15)$, then \mathcal{S} is the characteristic function of a solid.
- (e) If $(\Delta, n) = (8, 30)$, then \mathcal{S} is the sum of the characteristic functions of two disjoint solids.

Theoretical proofs and generalizations can e.g. be found in [Kur21, Section 11] or [HKK18]. For $q > 2$ a much stronger result is known. [GS03b, Theorem 13] directly implies:

Theorem 7. Let \mathcal{M} be a q^r -divisible multiset of cardinality $\delta \cdot \frac{q^{r+1}-1}{q-1}$ over \mathbb{F}_q , where $r \in \mathbb{N}_{>0}$. If $q > 2$ and $1 \leq \delta < \varepsilon$, where $q + \varepsilon$ is the size of the smallest non-trivial blocking sets in $\text{PG}(2, q)$, then there exist $(r+1)$ -spaces S_1, \dots, S_δ such that

$$\mathcal{M} = \sum_{i=1}^{\delta} \chi_{S_i},$$

i.e., \mathcal{M} is the sum of $(r+1)$ -spaces.

Using dimension arguments, c.f. the proof of Lemma 15, we conclude from Lemma 6.(c)-(e):

Corollary 8. *Let \mathcal{P} be a vector space partition of $\text{PG}(v-1, 2)$, then \mathcal{P} cannot have a supertail of one of the following types:*

- $2^3 1^5$;
- $3^1 2^1 1^5$; and
- $3^2 2^1 1^{13}$, $3^2 2^2 1^{10}$, $3^2 2^3 1^7$, $3^2 2^4 1^4$, $3^1 2^6 1^5$.

Lemma 6.(a) yields the fact that every vector space partition of $\text{PG}(v-1, 2)$ of type $(v-1)^{m_{v-1}} \dots 2^{m_2} 1^3$ is reducible (assuming $v > 2$), i.e., the three points have to form a line. Similarly, from Lemma 6.(b) we can conclude that every vector space partition of $\text{PG}(v-1, 2)$ of type $(v-1)^{m_{v-1}} \dots 3^{m_3} 1^7$ is reducible (assuming $v > 3$), i.e., the seven points have to form a plane.

While Lemma 6 discusses the parameters of (repeated) simplex codes, i.e., duals of Hamming codes, there are also known uniqueness results for the parameters of first order Reed-Muller codes, see e.g. [KM21]. Translated to geometry we have:

Lemma 9. *Let \mathcal{S} be a 2^r -divisible set of 2^{r+1} points in $\text{PG}(v-1, 2)$, where $r \in \mathbb{N}_{>0}$. Then, \mathcal{S} is the characteristic function of an affine $(r+1)$ -space.*

In our context an important implication is that such a set \mathcal{S} does not contain a line, so that we conclude:

Lemma 10. *Let \mathcal{P} be a vector space partition of $\text{PG}(v-1, 2)$, then \mathcal{P} cannot have a supertail of one of the following types:*

- $2^1 1^5$ and
- $3^0 2^1 1^{13}$, $3^0 2^2 1^{10}$, $3^0 2^3 1^7$, $3^0 2^4 1^4$.

For larger field sizes there exist examples different to affine subspaces. They can be obtained by the so-called cylinder construction, see e.g. [DBDMS19], and share the property that those point sets also do not contain a line. A few parameters where q^r -divisible sets of q^{r+1} points in $\text{PG}(n-1, q)$ have to be obtained by the cylinder construction are recently discussed in [KM21].

We say that a multiset of points \mathcal{M} in $\text{PG}(v-1, q)$ is *spanning* if the points P with non-zero multiplicity $\mathcal{M}(P) > 0$ span the entire ambient space. For a given multiset \mathcal{M} of points in $\text{PG}(v-1, q)$ we denote by a_i the number of hyperplanes H such that $\mathcal{M}(H) = i$. The vector $(a_i)_{i \in \mathbb{N}}$ is called the *spectrum* of \mathcal{M} . If \mathcal{M} is spanning, then we have $a_{\#\mathcal{M}} = 0$. By considering \mathcal{M} restricted to $K \cong \text{PG}(k-1, q)$ we can always assume that \mathcal{M} is spanning if we choose a suitable integer for k . For the ease of notation we assume that \mathcal{M} is spanning in $\text{PG}(k-1, q)$ in the following. Counting the number of hyperplanes in $\text{PG}(k-1, q)$ gives

$$\sum_i a_i = \frac{q^k - 1}{q - 1} \quad (4)$$

and counting the number of pairs of points and hyperplanes gives

$$\sum_i i a_i = \#\mathcal{M} \cdot \frac{q^{k-1} - 1}{q - 1}. \quad (5)$$

For the third equation we assume $\mathcal{M}(P) \in \{0, 1\}$ for every point P , i.e., there is no point with multiplicity at least 2. Double-counting the incidences between pairs of elements in \mathcal{M} and hyperplanes gives

$$\sum_i \binom{i}{2} a_i = \binom{\#\mathcal{M}}{2} \cdot \frac{q^{k-2} - 1}{q - 1}. \quad (6)$$

We call the equations (4)-(6) the *standard equations* for sets of points.

3 Vector space partitions in $\text{PG}(7, 2)$

For each dimension v of the ambient space and each field size q the packing condition in Equation (1) combined with the dimension condition in Equation (2) leave over a finite list of possibly types of vector space partitions in $\text{PG}(v-1, q)$. The conditions on the supertail mentioned in the previous section eliminate a few more cases. Here we will treat the case of $\text{PG}(7, 2)$, which is quite comprehensive compared to the case $\text{PG}(6, 2)$. In the end it will turn out that there are more than ten thousand different possible types of vector space partitions in $\text{PG}(7, 2)$. E.g. there are vector space partitions of type $4^4 3^{12} 2^{61-i} 1^{5+3i}$ for each integer $0 \leq i \leq 61$. Of course it is sufficient to give constructions for the “irreducible cases” only. I.e., in our example it suffice to give an example of type $4^4 3^{12} 2^{61} 1^5$ and then to replace i lines by its three contained points each. We have utilized the following general reduction rules:

Lemma 11. *Let \mathcal{P} be a vector space partition of type $(v-1)^{m_{v-1}} \dots 2^{m_2} 1^{m_1}$ in $\text{PG}(v-1, 2)$.*

- *If $m_2 > 0$, then there also exists a vector space partition of type $(v-1)^{m_{v-1}} \dots 3^{m_3} 2^{m_2-1} 1^{m_1+3}$.*
- *If $m_3 > 0$, then there also exists a vector space partition of type $(v-1)^{m_{v-1}} \dots 4^{m_4} 3^{m_3-1} 2^{m_2} 1^{m_1+7}$.*
- *If $m_3 > 0$, then there also exists a vector space partition of type $(v-1)^{m_{v-1}} \dots 4^{m_4} 3^{m_3-1} 2^{m_2+1} 1^{m_1+4}$.*
- *If $m_4 > 0$, then there also exists a vector space partition of type $(v-1)^{m_{v-1}} \dots 5^{m_5} 4^{m_4-1} 3^{m_3} 2^{m_2} 1^{m_1+15}$.*
- *If $m_4 > 0$, then there also exists a vector space partition of type $(v-1)^{m_{v-1}} \dots 5^{m_5} 4^{m_4-1} 3^{m_3} 2^{m_2+5} 1^{m_1}$.*
- *If $m_4 > 0$, then there also exists a vector space partition of type $(v-1)^{m_{v-1}} \dots 5^{m_5} 4^{m_4-1} 3^{m_3+1} 2^{m_2} 1^{m_1+8}$.*

Nevertheless, still a lot of examples of vector space partitions need to be constructed. To this end we utilize integer linear programming (ILP) formulations. For each i -space S we introduce the binary variable $x_S^i \in \{0, 1\}$ with the meaning $x_S^i = 1$ iff S is contained in the vector space partition \mathcal{P} in $\text{PG}(v-1, q)$. The partitioning condition is modeled by

$$\sum_{i=1}^{v-1} \sum_{S: \dim(S)=i, P \leq S} x_S^i = 1 \quad (7)$$

for each point P . Introducing the counting variables

$$m_i = \sum_{S: \dim(S)=i} x_S^i \quad (8)$$

for each dimension $1 \leq i \leq v-1$, we can prescribe the type or maximizing/minimizing certain values m_i while prescribing the others. Of course this general ILP has a huge number of variables and constraints. E.g. there are 200 787 solids in $\text{PG}(7, 2)$. Luckily enough we can use the collineation group of $\text{PG}(7, 2)$ to reduce the search space a bit. It is well known that the collineation group acts transitively on i -spaces as well as pairs of disjoint i - and j -spaces, including the case $i = j$. For triples of disjoint i -spaces A, B, C the orbits have $\dim(\langle A, B, C \rangle)$ as invariant that can vary between $2i$ and $\min\{v, 3i\}$, where $v = 8$ in our situation. So, we can prescribe up to three elements of the vector space partition and search for the others.

In the special case where m_4 is relatively large, we have considered a Desarguesian spread for solids in $\text{PG}(7, 2)$, see e.g. [Lun99]. So, let \mathcal{S} be such a set of 17 pairwise disjoint solids in $\text{PG}(7, 2)$ (that form a vector space partition of type 4^{17}). We can restrict the search space for the ILP by forcing the used solids to be contained in \mathcal{S} , i.e., we set $x_S^4 = 0$ for every solid S that is not contained in \mathcal{S} . Another starting point for vector space partitions are so-called lifted MRD (maximum rank distance) codes, see e.g. [SSW19], that e.g. give vector space partitions of types $6^1 2^{64}$, $5^1 3^{32}$, and also 4^{17} .

For ILP formulations of search problems in incidence structures there is a general (heuristic) method to reduce the search space – the so-called Kramer–Mesner method [KM76]. Here a suitable subgroup G of the automorphism group of the desired object is assumed. The action of the group G partitions the variables

into orbits and we assume that variables in the same orbit have the same value. I.e., in our context the vector space partition consists of entire orbits of solids, planes, and so on. This approach reduces the number of variables in terms of the order of G . Also the number of constraints is automatically reduced since usually several constraints become identical if we use orbit variables. For a more detailed and precise description of the method we refer e.g. to [KK08] where sets of subspaces in $\text{PG}(v-1, q)$ with restricted intersection dimension were considered. Via this method we can, of course, find examples whose automorphism group contains G as a subgroup only. So, as mentioned, the method is heuristic, but quite successfully applied in the literature. In our context we have used groups G with orders up to 16.

As our setting is quite comprehensive we refrain from stating the explicit details for the cases. However, explicit vector space partitions for specific types in $\text{PG}(7, 2)$ can be obtained from the author upon request. While it took quite some time to compute an example for each feasible case, we now focus on the complementary non-existence results.

Lemma 12. *Let \mathcal{P} be a vector space partition of $\text{PG}(v-1, 2)$, then \mathcal{P} cannot have a supertail of type 2^{415} .*

Proof. There are three different 4-divisible sets of cardinality 17, one for each of the dimensions $k \in \{6, 7, 8\}$, see [HHK⁺17]. Only the 6-dimensional point set admits three disjoint lines and there cannot be four disjoint lines. \square

As a consequence, there does not exist a vector space partition of type $3^{34}2^{415}$ of $\text{PG}(7, 2)$ but there exist vector space partitions of type $3^{34}2^{3-i}1^{8+3i}$ for all integers $0 \leq i \leq 3$. A set of 34 pairwise disjoint planes in $\text{PG}(7, 2)$ was constructed for the first time in [EZJS⁺10]. Now several thousand non-isomorphic examples are known, see [HKK18].

Lemma 13. *For $k \geq 3$ let K_1, K_2, K_3 be three pairwise disjoint k -spaces in $\text{PG}(v-1, 2)$ and \mathcal{L} be the set of lines that intersect each K_i in exactly a point. Then, the lines in \mathcal{L} are pairwise disjoint and $\{L \cap K_i : L \in \mathcal{L}\}$ forms a subspace for each index $1 \leq i \leq 3$.*

Proof. First we show that the lines in \mathcal{L} are pairwise disjoint. To the contrary we assume that \mathcal{L} contains two lines $L_1 = \langle A, B \rangle$ and $L_2 = \langle A, B' \rangle$, where we assume w.l.o.g. that $A \leq K_1$ and $B, B' \leq K_2$. Then $A + B, A + B' \leq K_3$, so that $B + B' \in K_2 \cap K_3$, which is a contradiction.

For the second part we show the statement for K_1 and assume that $L_1 = \langle A_1, B_1 \rangle$ and $L_2 = \langle A_2, B_2 \rangle$ are two different lines in \mathcal{L} with $A_1, A_2 \leq K_1$ and $B_1, B_2 \leq K_2$. Then also $\langle A_1 + A_2, B_1 + B_2 \rangle$ is in \mathcal{L} since $A_1 + A_2 \leq K_1$, $B_1 + B_2 \leq K_2$, and $A_1 + A_2 + B_1 + B_2 \leq K_3$. \square

Lemma 14. *For $k \geq 3$ let K_1, K_2, K_3 be three pairwise disjoint k -spaces in $\text{PG}(2k-1, 2)$ and \mathcal{L} be the set of lines that intersect each K_i in exactly a point. Then, $\#\mathcal{L} = 2^k - 1$ and the pairwise disjoint lines in \mathcal{L} partition the point set $K_1 \cup K_2 \cup K_3$.*

Proof. We have already mentioned that there is a unique isomorphism type under the operation of the collineation group of $\text{PG}(2k-1, 2)$ since $\dim(\langle K_i, K_j \rangle) = 2k$ for all $1 \leq i < j \leq 3$. W.l.o.g. we choose $K_1 = \langle e_1, \dots, e_k \rangle$, $K_2 = \langle e_{k+1}, \dots, e_{2k} \rangle$, and $K_3 = \langle e_1 + e_{k+1}, \dots, e_k + e_{2k} \rangle$, where e_i denotes the i th unit vector. Let $P_1 = \langle \sum_{i=1}^k a_i e_i \rangle$ be the unique point of such a line in K_1 and $P_2 = \langle \sum_{i=1}^k b_i e_{i+k} \rangle$ be the unique intersection of the line with K_2 , where $a_1, \dots, a_k, b_1, \dots, b_k \in \{0, 1\}$. Since the third point of the line is given by $P_3 = \sum_{i=1}^k (a_i e_i + b_i e_{i+k}) \in K_3$, we conclude $a_i = b_i$ for all $1 \leq i \leq k$. Moreover, we have to exclude the case $a_1 = \dots = a_k$, so that $2^k - 1$ possibilities remain. \square

Lemma 15. *Let \mathcal{P} be a vector space partition of $\text{PG}(v-1, 2)$, then \mathcal{P} cannot have a supertail of one of the following types: $3^4 2^{415}$, $3^4 2^3 1^8$, $3^4 2^2 1^{11}$, $3^4 2^1 1^{14}$, $3^4 2^0 1^{17}$, and $3^1 2^{11} 1^5$.*

Proof. The stated types for supertails all consist of 45 points. The 8-divisible sets \mathcal{S} of cardinality 45 in $\text{PG}(v-1, 2)$ have been computationally classified in [HHK⁺17]. Either \mathcal{S} is the sum of the characteristic functions of three pairwise disjoint solids or \mathcal{S} is isomorphic to the points P_1, \dots, P_9 of a projective base in $\text{PG}(7, 2)$ and the $\binom{9}{2} = 36$ remaining points on the lines $P_i P_j$.

We can easily check that the latter case does not contain a plane in its support. Thus, we can assume that $\mathcal{S} = \chi_{S_1} + \chi_{S_2} + \chi_{S_3}$, where S_1, S_2, S_3 are pairwise disjoint solids. Each plane E contained in the support of \mathcal{S} has to intersect each of the solids S_i in a d_i -dimensional subspace where $d_i \in \{0, 1, 2, 3\}$. Since two lines contained in a plane have to intersect in a point, the only possibility is that one of the solids S_i completely contains the plane E . However, this is impossible for four pairwise disjoint planes. So, let us finally assume that \mathcal{S} has type $3^1 2^{11} 1^5$ and that the unique plane E is contained in S_1 . The five points form a 2-divisible set, so that they have to be isomorphic to a projective base \mathcal{B} of size 5, see e.g. [HHK⁺17]. So, at most four of these five points can be contained in S_1 . Denoting the set of lines that intersect S_1, S_2 , and S_3 in exactly a point by \mathcal{L} and setting $\alpha := \#\mathcal{L}$, we observe $\alpha \geq 4$. So, S_2 and S_3 can contain at most three lines each and $\alpha \geq 5$. Since \mathcal{B} is 2-divisible we have $\alpha \neq 8$, so that $5 \leq \alpha \leq 7$.

Note that no three points in $\mathcal{P}_1 := \{L \cap S_1 : L \in \mathcal{L}\}$ can form a line, so that Lemma 13 yields that also no three points in $\mathcal{P}_i := \{L \cap S_i : L \in \mathcal{L}\}$ can form a line for all $1 \leq i \leq 3$. Thus the $\alpha \geq 5$ points in \mathcal{P}_3 have to span a solid that is also contained in $\langle S_1, S_2 \rangle$, so that we conclude $n = 8$ for the dimension of the ambient space. Note that for $i = 2, 3$ the points of S_i are partitioned by the lines contained in S_i and the points $\mathcal{P}_i \cup (\mathcal{B} \cap S_i)$. Note that the points in $\mathcal{P}_i \cup (\mathcal{B} \cap S_i)$ form a 2-divisible set in $\text{PG}(3, 2)$ whose cardinality is a multiple of three. For cardinality three we can apply Lemma 6 to conclude that those points form a line. If $\#(\mathcal{P}_i \cup (\mathcal{B} \cap S_i)) = 3\beta$, where $2 \leq \beta \leq 5$, then there exists a unique 2-divisible set of 3β points in $\text{PG}(3, 2)$, see [HHK⁺17]. In all cases the span of the points has dimension 3 and an example is given by β disjoint lines. Thus, for $i = 2, 3$ the points in $\mathcal{P}_i \cup (\mathcal{B} \cap S_i)$ can be partitioned into lines, so that $\alpha = \#\mathcal{P}_i \leq 2\beta = \frac{2}{3} \cdot (\alpha + \#(\mathcal{B} \cap S_i))$, which is equivalent to $\alpha \leq 2\#(\mathcal{B} \cap S_i)$. Since $\#(\mathcal{B} \cap S_2) + \#(\mathcal{B} \cap S_3) = \#\mathcal{B} = 5$ and $\alpha \geq 5$ this is impossible. \square

So, especially there does not exist a vector space partition of type $4^{14} 3^4 1^{17}$ or $4^{14} 3^1 2^{11} 1^5$ in $\text{PG}(7, 2)$. (We remark that we have also checked the more involved second part of the proof of Lemma 15, i.e., that the set of points of three disjoint solids cannot be partitioned in a plane, eleven lines, and five points computationally using an ILP formulation.)

Lemma 16. *In $\text{PG}(7, 2)$ no vector space partitions of types $4^{13} 3^6 2^{6-i} 1^{3i}$ for $0 \leq i \leq 6$, types $4^{13} 3^5 2^{7-i} 1^{4+3i}$ for $0 \leq i \leq 7$, and type $4^{13} 3^4 2^9 1^5$ exist.*

Proof. Assume that \mathcal{P} is a vector space partition of one of those types and observe that the set \mathcal{H} of points that are not covered by the 13 solids of \mathcal{P} forms an 8-divisible set of 60 points. Let k be the dimension of the span of \mathcal{H} . For the ease of notation we assume that \mathcal{H} is embedded in $\text{PG}(k-1, 2)$ and denote by a_i the number of hyperplanes containing exactly i points from \mathcal{H} , so that the standard equations are given by

$$\sum_{i=0}^6 a_{4+8i} = 2^k - 1, \quad (9)$$

$$\sum_{i=0}^6 (4+8i) \cdot a_{4+8i} = 60 \cdot (2^{k-1} - 1), \text{ and} \quad (10)$$

$$\sum_{i=0}^6 (4+8i)(3+8i) \cdot a_{4+8i} = 60 \cdot 59 \cdot (2^{k-2} - 1). \quad (11)$$

Using Equation (9) and Equation (10) we conclude

$$\sum_{i=0}^6 i \cdot a_{4+8i} = 13 \cdot 2^{k-2} - 7 \quad (12)$$

Equation (11) minus 56 times Equation (10) minus 12 times Equation (9) gives

$$\sum_{i=0}^6 i^2 \cdot a_{4+8i} = 691 \cdot 2^{k-6} - 49 \quad (13)$$

after a final division by 64. Now, Equation (13) minus seven times Equation (12) plus twelve times Equation (9) yields

$$\sum_{i=0}^6 (i-3)(i-4)a_{4+8i} = 3 \cdot 2^{k-6} - 12. \quad (14)$$

Since $(i-3)(i-4) \geq 0$ and $a_{4+8i} \geq 0$ for all i , we have $k \geq 8$. Due to the ambient space $\text{PG}(7, 2)$ for \mathcal{P} we are only interested in the case $k = 8$ where $\sum_{i=0}^7 (i-3)(i-4)a_{4+8i} = 0$. Thus, the unique solution is given by $a_{36} = 60$, $a_{28} = 195$, and $a_i = 0$ otherwise.

We have used `LinCode`, see [BBK21], to enumerate all projective $[60, 8, \{24, 32\}]_2$ codes. There are exactly 12 non-isomorphic such codes, c.f. [BFWW06]. We have computationally checked that the corresponding sets of points contain at most five disjoint planes. Five disjoint planes can be obtained in just one of the twelve cases. However, we have computationally checked that in this case no 13 pairwise disjoint solids can be chosen outside this point set. Four planes and nine lines that are pairwise disjoint can occur in two of the twelve cases. Again, we have computationally checked that in those two cases no 13 pairwise disjoint solids can be chosen outside the point set. \square

Lemma 17. *If \mathcal{P} is a vector space partition of type $4^{12}3^81^{19}$ in $\text{PG}(7, 2)$, then the point set of the points in the subspaces of dimension at most 3 is isomorphic to the columns of*

$$\begin{pmatrix} 1111111111111111111111111111111100000000000000000000000000000010000000 \\ 0000000000000011111111111111111111111111110000000000000000000001000000 \\ 000000011111110000000011111110000000111111111111110000000000000001000000 \\ 001111000011110000111100001111000111100011111001111110000000000000010000 \\ 0010011001100010001001100000111011000100010011001001111111111100000001000 \\ 010010100010000001001010111001101110001101000110100001000111111000000100 \\ 11011101010110110011110011000100001100110010110100110110110011011100000010 \\ 00010010001001001011001000110111010101000110001000101110101011010000001 \end{pmatrix},$$

$$\begin{pmatrix} 1111111111111111111111111111111100000000000000000000000000000000000001000000 \\ 0000000000000001111111111111111111111111111000000000000000000000000001000000 \\ 00000001111111000000011111110000000111111111111110000000000000000001000000 \\ 0001111000011110000111100001111000111100011111110000000000000000010000 \\ 01100000011001100110011001100000110011000000110110011011111111000000001000 \\ 1010000010101010101010101000010101010000010110101011010001111111000000100 \\ 011001100111010100110010101001100010100011000001100001100110011011100000010 \\ 101010100000101011000000110010101100110101000011001010110101011010000001 \end{pmatrix},$$

or

$$\begin{pmatrix} 111111111111111111111111111111110000000000000000000000000000000000000001000000 \\ 000000000000000111111111111111111111111111100000000000000000000000001000000 \\ 00000001111111000000001111111000000011111111111111000000000000000001000000 \\ 00011110000111100001111000011110001111000111110001111110000000000000010000 \\ 0110000001100110011001100110000011001100000011011001101111111000000001000 \\ 1010000010101010101010101000010101010000010110101011010001111110000001000 \\ 011001100111010100110010101001100010100011000001100001100110011011100000010 \\ 110010101010010101010100010100000110101111110111101101010110110000001 \end{pmatrix}.$$

Proof. Observe that the set \mathcal{H} of points that are not covered by the 12 solids of \mathcal{P} forms an 8-divisible set of 75 points. Let k be the dimension of the span of \mathcal{H} . For the ease of notation we assume that \mathcal{H} is embedded in $\text{PG}(k-1, 2)$ and denote by a_i the number of hyperplanes containing exactly i points from \mathcal{H} . Similar as in the proof of Lemma 16 we use the so standard equations $\sum_{i=0}^8 a_{3+8i} = 2^k - 1$, $\sum_{i=0}^8 (3+8i) \cdot a_{3+8i} = 75 \cdot (2^{k-1} - 1)$, and $\sum_{i=0}^8 (3+8i)(2+8i) \cdot a_{3+8i} = 75 \cdot 74 \cdot (2^{k-2} - 1)$ to conclude

$$\sum_{i=0}^8 i \cdot a_{3+8i} = 69 \cdot 2^{k-4} - 9,$$

$$\sum_{i=0}^8 i^2 \cdot a_{3+8i} = 1209 \cdot 2^{k-6} - 81, \text{ and}$$

$$\sum_{i=0}^8 (i-4)(i-5)a_{3+8i} = 4 \cdot 2^{k-6} - 20.$$

Since $(i-4)(i-5) \geq 0$ and $a_{3+8i} \geq 0$ for all i , we have $k \geq 8$. Due to the ambient space $\text{PG}(7, 2)$ for \mathcal{P} we are only interested in the case $k = 8$ where $\sum_{i=0}^8 (i-4)(i-5)a_{3+8i} = 0$. Thus, the unique solution is given by $a_{43} = 75$, $a_{35} = 180$, and $a_i = 0$ otherwise.

Using the notation $[n, k]_2$ -code for a binary k -dimensional code of effective length n , \mathcal{H} is given by the columns of a projective $[75, 8]_2$ -two-weight code C with weights 32 and 40. Since \mathcal{H} contains a plane C can be obtained recursively by lengthening $[74, 7]_2$, $[72, 6]_2$, and $[68, 5]_2$ codes with maximum possible column multiplicities at most 2, 4, and 8, respectively, where all non-zero weights are contained in $\{32, 40\}$. Using `LinCode`, see [BBK21], we enumerated 38 $[68, 5]_2$, 4286 $[72, 6]_2$, 245736 $[74, 7]_2$, and 9964 $[75, 8]_2$ codes. The complete enumeration took 85 hours of computation time. Checking which of those 9964 point sets allow eight disjoint planes leaves just seven possibilities. Using an ILP formulation for a partition of the complement into 12 solids leaves just the three mentioned cases. \square

Lemma 18. *In $\text{PG}(7, 2)$ no vector space partition of type $4^{12}3^82^{5-i}1^{4+3i}$ for $2 \leq i \leq 5$ exists.*

Proof. Assume that \mathcal{P} is a vector space partition of one of those types. Replacing the lines by their contained three points gives a vector space partition \mathcal{P}' of type $4^{12}3^81^{19}$, so that we can apply Lemma 17. Let \mathcal{H} be the corresponding point set of the points in the subspaces of dimension at most 3. Using an integer linear programming formulation we have checked that we can pack at most one line if we also pack eight disjoint planes into \mathcal{H} . \square

We remark that from the three 4-divisible projective binary linear codes of length 19, see e.g. [HHK⁺17], one contains five disjoint lines and the other two contain no pair of disjoint lines.

Lemma 19. *Let \mathcal{S} be a 4-divisible multiset of points of cardinality 20 and dimension at most 8 that contains five disjoint lines, then, up to symmetry, \mathcal{S} is given by the columns of*

$$\begin{pmatrix} 10000011101000010110 \\ 01000000011101101100 \\ 00100011110110000010 \\ 00010010011010001110 \\ 00001001100001111100 \\ 00000101111001001010 \\ 00000000110000000101 \end{pmatrix}.$$

Moreover, we have $\dim(\mathcal{S}) = 7$ and the spectrum is given by $(a_8, a_{12}, a_{16}) = (67, 59, 1)$.

Proof. The projective 4-divisible binary linear codes of cardinality 20 have been classified in [HHK⁺17]. Their counts per dimension are given by 7^28^{49} . By a direct enumeration we have checked which corresponding point sets contain five disjoint lines. \square

Lemma 20. *In $\text{PG}(7, 2)$ no vector space partition of type $4^43^{25}2^{515}$ exists.*

Proof. Assume that \mathcal{P} is a vector space partition of this type and observe that the set \mathcal{H} of points that are not covered by the 4 solids and the 25 planes of \mathcal{P} forms an 4-divisible set of 20 points that contains five disjoint lines. The unique possibility up to symmetry is determined in Lemma 19. It turns out that the ILP formulation for a vector space partition of type $4^43^{25}2^{515}$ is infeasible when prescribing these 20 points. \square

Lemma 21. *In $\text{PG}(7, 2)$ no vector space partition of type $4^{11}3^{10}2^{515}$ exists.*

Proof. Assume that \mathcal{P} is a vector space partition of type $4^{11}3^{10}2^{515}$ and observe that the set \mathcal{H} of points that are not covered by the 11 solids of \mathcal{P} forms an 8-divisible set of 90 points. Let k be the dimension of the span of \mathcal{H} . For the ease of notation we assume that \mathcal{H} is embedded in $\text{PG}(k-1, 2)$ and denote by a_i the number of hyperplanes containing exactly i points from \mathcal{H} . Similar as in the proof of Lemma 16 we use the standard equations $\sum_{i=0}^{11} a_{2+8i} = 2^k - 1$, $\sum_{i=0}^{11} (2+8i) \cdot a_{2+8i} = 90 \cdot (2^{k-1} - 1)$, and $\sum_{i=0}^{11} (2+8i)(1+8i) \cdot a_{2+8i} =$

$90 \cdot 89 \cdot (2^{k-2} - 1)$ to conclude

$$\begin{aligned} \sum_{i=0}^{11} i \cdot a_{2+8i} &= 43 \cdot 2^{k-3} - 11, \\ \sum_{i=0}^{11} i^2 \cdot a_{2+8i} &= 3743 \cdot 2^{k-7} - 121, \text{ and} \\ \sum_{i=0}^{11} (i-5)(i-6)a_{3+8i} &= 15 \cdot 2^{k-7} - 30. \end{aligned}$$

Since $(i-5)(i-6) \geq 0$ and $a_{3+8i} \geq 0$ for all i , we have $k \geq 8$. Due to the ambient space $\text{PG}(7, 2)$ for \mathcal{P} we are only interested in the case $k = 8$ where $\sum_{i=0}^{11} (i-5)(i-6)a_{3+8i} = 0$. Thus, the unique solution is given by $a_{50} = 90$, $a_{42} = 165$, and $a_i = 0$ otherwise.

If \mathcal{H} contains a solid S in its support, then there are 75 points in \mathcal{H} that are not contained in S and each hyperplane contain either 35 or 43 points of these. Via projective $[75, 8]_2$ -two-weight codes C with weights 32 and 40 such point sets have already been enumerated in Lemma 18. Via ILP computations we have filtered out which of the corresponding point sets can be completed by 12 solids to a vector space partition of $\text{PG}(7, 2)$. Only 42 codes remain and we extended them in all possible ways by a four-dimensional simplex code such that the code remains projective. After filtering out isomorphic codes we have again used an ILP formulation to check which point sets can be completed by 11 solids to a vector space partition of $\text{PG}(7, 2)$. For the remaining 245 point sets we have checked which allow to pack 10 disjoint planes into them via ILP computations. This was possible in 10 cases only and we finally have checked using ILP computations that we can pack at most 4 disjoint lines when we also pack 10 disjoint planes into the point set. In other words, the 20 points that are not covered by the 11 solids and the 10 planes do not form a 4-divisible point set as specified in Lemma 19.

These computations show that in the remaining part we can assume that \mathcal{H} does not contain a full solid in its support. Now we have enumerated the possibilities of four pairwise disjoint solids in $\text{PG}(7, 2)$ up to symmetry. By exhaustive enumeration we have extended the four prescribed solids to 11 solids in total that are pairwise disjoint and discarded all cases that allow the addition of a 12th such solid. For all of these cases we have exhaustively enumerated all vector space partitions of type $4^{11}3^{10}1^{20}$ and determined the maximum number of disjoint lines that we can pack into the remaining 20 points. In all cases the answer was at most 4. \square

We remark that we have also tried to enumerate the projective $[90, 8]_2$ -two-weight codes C with weights 40 and 48 directly. However, we stopped the computations after having reached more than 1.5 million different codes.

For every type satisfying the numerical conditions of Equation (1) and Equation (2) that is not excluded by one of the previous lemmas there indeed exists a corresponding vector space partition of $\text{PG}(7, 2)$. We summarize the set of feasible parameters as follows:

Theorem 22. *Let \mathcal{P} be a vector space partition of $\text{PG}(7, 2)$, then \mathcal{P} has one of the following types:*

- $7^2 1^{128}$;
- $6^2 2^{64-i} 1^{3i}$, where $0 \leq i \leq 64$;
- $5^1 3^{32}$;
- $5^1 3^{31-3j} 2^{1-i+7j} 1^{4+3i}$, where $0 \leq i \leq 1+7j$ and $0 \leq j \leq 10$;
- $5^1 3^{29-3j} 2^{7-i+7j} 1^{3i}$, where $0 \leq i \leq 7+7j$ and $0 \leq j \leq 9$;
- $5^1 3^{27-3j} 2^{10-i+7j} 1^{5+3i}$, where $0 \leq i \leq 10+7j$ and $0 \leq j \leq 9$;

- 4^{17} ;
- $4^{16}3^{11}8$;
- $4^{16}2^{5-i}1^{3i}$, where $0 \leq i \leq 5$;
- $4^{15}3^21^{16}$;
- $4^{15}3^12^{5-i}1^{8+3i}$, where $0 \leq i \leq 5$;
- $4^{15}2^{10-i}1^{3i}$, where $0 \leq i \leq 10$;
- $4^{14}3^{3-3j}2^{8-i+7j}1^{3i}$, where $0 \leq i \leq 8+7j$ and $0 \leq j \leq 1$;
- $4^{14}3^22^{9-i}1^{4+3i}$, where $0 \leq i \leq 9$;
- $4^{14}3^12^{10-i}1^{8+3i}$, where $0 \leq i \leq 10$;
- $4^{13}3^42^{8-i}1^{8+3i}$, where $0 \leq i \leq 8$;
- $4^{13}3^{3-3j}2^{13-i+7j}1^{3i}$, where $0 \leq i \leq 13+7j$ and $0 \leq j \leq 1$;
- $4^{13}3^22^{14-i}1^{4+3i}$, where $0 \leq i \leq 14$;
- $4^{13}3^12^{16-i}1^{5+3i}$ where $0 \leq i \leq 16$;
- $4^{12}3^82^{1-i}1^{16+3i}$, where $0 \leq i \leq 1$;
- $4^{12}3^{7-3j}2^{7-i+7j}1^{5+3i}$, where $0 \leq i \leq 7+7j$ and $0 \leq j \leq 2$;
- $4^{12}3^{6-3j}2^{11-i+7j}1^{3i}$, where $0 \leq i \leq 11+7j$ and $0 \leq j \leq 2$;
- $4^{12}3^{5-3j}2^{12-i+7j}1^{4+3i}$, where $0 \leq i \leq 11+7j$ and $0 \leq j \leq 1$;
- $4^{11}3^102^{4-i}1^{8+3i}$, where $0 \leq i \leq 4$;
- $4^{11}3^9-3j2^{9-i+7j}1^{3i}$, where $0 \leq i \leq 9+7j$ and $0 \leq j \leq 3$;
- $4^{11}3^{8-3j}2^{10-i+7j}1^{4+3i}$, where $0 \leq i \leq 10+7j$ and $0 \leq j \leq 2$;
- $4^{11}3^{7-3j}2^{12-i+7j}1^{5+3i}$, where $0 \leq i \leq 12+7j$ and $0 \leq j \leq 2$;
- $4^{10}3^{15}$;
- $4^{10}3^{14-3j}2^{1-i+7j}1^{4+3i}$, where $0 \leq i \leq 1+7j$ and $0 \leq j \leq 4$;
- $4^{10}3^{13}2^{2-i}1^{8+3i}$, where $0 \leq i \leq 2$;
- $4^{10}3^{12-3j}2^{7-i+7j}1^{3i}$, where $0 \leq i \leq 7+7j$ and $0 \leq j \leq 4$;
- $4^{10}3^{10-3j}2^{10-i+7j}1^{5+3i}$, where $0 \leq i \leq 10+7j$ and $0 \leq j \leq 3$;
- $4^93^{16}1^8$;
- $4^93^{15-3j}2^{5-i+7j}1^{3i}$, where $0 \leq i \leq 5+7j$ and $0 \leq j \leq 5$;
- $4^93^{14-3j}2^{6-i+7j}1^{4+3i}$, where $0 \leq i \leq 6+7j$ and $0 \leq j \leq 4$;
- $4^93^{13-3j}2^{8-i+7j}1^{5+3i}$, where $0 \leq i \leq 8+7j$ and $0 \leq j \leq 4$;
- $4^83^{17-3j}2^{4-i+7j}1^{4+3i}$, where $0 \leq i \leq 4+7j$ and $0 \leq j \leq 5$;
- $4^83^{16-3j}2^{6-i+7j}1^{5+3i}$, where $0 \leq i \leq 6+7j$ and $0 \leq j \leq 5$;

- $4^8 3^{15-3j} 2^{10-i+7j} 1^{3i}$, where $0 \leq i \leq 10 + 7j$ and $0 \leq j \leq 5$;
- $4^7 3^{19} 2^{3-i} 1^{8+3i}$, where $0 \leq i \leq 3$;
- $4^7 3^{18-3j} 2^{8-i+7j} 1^{3i}$, where $0 \leq i \leq 8 + 7j$ and $0 \leq j \leq 6$;
- $4^7 3^{17-3j} 2^{9-i+7j} 1^{4+3i}$, where $0 \leq i \leq 9 + 7j$ and $0 \leq j \leq 5$;
- $4^7 3^{16-3j} 2^{11-i+7j} 1^{5+3i}$, where $0 \leq i \leq 11 + 7j$ and $0 \leq j \leq 5$;
- $4^6 3^{21-3j} 2^{6-i+7j} 1^{3i}$, where $0 \leq i \leq 6 + 7j$ and $0 \leq j \leq 7$;
- $4^6 3^{20-3j} 2^{7-i+7j} 1^{4+3i}$, where $0 \leq i \leq 7 + 7j$ and $0 \leq j \leq 6$;
- $4^6 3^{19-3j} 2^{9-i+7j} 1^{5+3i}$, where $0 \leq i \leq 9 + 7j$ and $0 \leq j \leq 6$;
- $4^5 3^{23-3j} 2^{5-i+7j} 1^{4+3i}$, where $0 \leq i \leq 5 + 7j$ and $0 \leq j \leq 7$;
- $4^5 3^{22-3j} 2^{7-i+7j} 1^{5+3i}$, where $0 \leq i \leq 7 + 7j$ and $0 \leq j \leq 7$;
- $4^5 3^{21-3j} 2^{11-i+7j} 1^{3i}$, where $0 \leq i \leq 11 + 7j$ and $0 \leq j \leq 7$;
- $4^4 3^{25} 2^{4-i} 1^{8+3i}$, where $0 \leq i \leq 4$;
- $4^4 3^{24-3j} 2^{9-i+7j} 1^{3i}$, where $0 \leq i \leq 9 + 7j$ and $0 \leq j \leq 8$;
- $4^4 3^{23-3j} 2^{10-i+7j} 1^{4+3i}$, where $0 \leq i \leq 10 + 7j$ and $0 \leq j \leq 7$;
- $4^4 3^{22-3j} 2^{12-i+7j} 1^{5+3i}$, where $0 \leq i \leq 12 + 7j$ and $0 \leq j \leq 7$;
- $4^3 3^{29} 2^{1-i} 1^4 + 3i$, where $0 \leq i \leq 1$;
- $4^3 3^{28} 2^{2-i} 1^{8+3i}$, where $0 \leq i \leq 2$;
- $4^3 3^{30-3j} 2^{7j-i} 1^{3i}$, where $0 \leq i \leq 7j$ and $0 \leq j \leq 10$;
- $4^3 3^{26-3j} 2^{8-i+7j} 1^{4+3i}$, where $0 \leq i \leq 8 + 7j$ and $0 \leq j \leq 8$;
- $4^3 3^{25-3j} 2^{10-i+7j} 1^{5+3i}$, where $0 \leq i \leq 10 + 7j$ and $0 \leq j \leq 8$;
- $4^2 3^{31} 1^8$;
- $4^2 3^{30-3j} 2^{5-i+7j} 1^{3i}$, where $0 \leq i \leq 5 + 7j$ and $0 \leq j \leq 10$;
- $4^2 3^{29-3j} 2^{6-i+7j} 1^{4+3i}$, where $0 \leq i \leq 6 + 7j$ and $0 \leq j \leq 9$;
- $4^2 3^{28-3j} 2^{8-i+7j} 1^{5+3i}$, where $0 \leq i \leq 8 + 7j$ and $0 \leq j \leq 9$;
- $4^1 3^{32-3j} 2^{4-i+7j} 1^{4+3i}$, where $0 \leq i \leq 4 + 7j$ and $0 \leq j \leq 10$;
- $4^1 3^{31-3j} 2^{6-i+7j} 1^{5+3i}$, where $0 \leq i \leq 6 + 7j$ and $0 \leq j \leq 10$;
- $4^1 3^{30-3j} 2^{10-i+7j} 1^{3i}$, where $0 \leq i \leq 10 + 7j$ and $0 \leq j \leq 10$;
- $4^0 3^{34} 2^{3-i} 1^{8+3i}$, where $0 \leq i \leq 3$;
- $4^0 3^{33-3j} 2^{8-i+7j} 1^{3i}$, where $0 \leq i \leq 8 + 7j$ and $0 \leq j \leq 11$;
- $4^0 3^{32-3j} 2^{9-i+7j} 1^{4+3i}$, where $0 \leq i \leq 9 + 7j$ and $0 \leq j \leq 10$;
- $4^0 3^{31-3j} 2^{11-i+7j} 1^{5+3i}$, where $0 \leq i \leq 11 + 7j$ and $0 \leq j \leq 10$;

In Proposition 27 in the appendix we also give a more explicit and extensive variant of the list of feasible types of vector space partitions of $\text{PG}(7, 2)$. A more compact variant is stated in Theorem 26.

4 Vector space partitions in $\text{PG}(v-1, q)$ for $v \leq 5$

The aim of this short section is to discuss the possible types of vector space partitions of $\text{PG}(v-1, q)$ for all dimensions $v \leq 5$ and arbitrary field size q . For $v = 1$ the ambient space consists of a single point itself, so that no vector space partition exists since we assume a maximum dimension of $v-1$ for the elements of a vector space partition. For the same reason all elements of a vector space partition of $\text{PG}(1, q)$ have dimension 1. I.e., the only possible type is 1^{m_1} where $m_1 = [2]_q = q+1$. For $v = 3$ the dimension condition yields that all elements of a vector space partition \mathcal{P} of $\text{PG}(2, q)$ have either dimension 1 or 2 and that dimension 2 can occur at most once. If \mathcal{P} does not contain an element of dimension 2, then its type is given by 1^{m_1} where $m_1 = [3]_q = q^2 + q + 1$. In that case \mathcal{P} is reducible since we may choose any line and choose it as an element of the vector space partition instead of its $q+1$ contained points. If \mathcal{P} is of type $2^1 1^{m_1}$, then we have $m_1 = [3]_q - [2]_q = q^2$.

Proposition 23. *The possible types of vector space partitions of $\text{PG}(3, q)$ are given by $2^{q^2+1-j} 1^{(q+1)j}$, where $0 \leq j \leq q^2 + 1$, and $3^1 1^{q^3}$.*

Proof. Directly implied by the packing and the dimension condition in equations (1) and (2). \square

All of the mentioned types can indeed be attained. A Desarguesian line spread in $\text{PG}(3, q)$ yields a vector space partition of type 2^{q^2+1} where we may replace arbitrary j lines by their contained points. Choosing an arbitrary plane in the ambient space leaves q^3 uncovered points. The latter vector space partition is irreducible. For vector space partitions of type $2^{q^2+1-j} 1^{(q+1)j}$ it is an interesting question which values of j do allow an irreducible vector space partition of that type. This problem is equivalent to the classification of the possible sizes of (inclusion) maximal partial line spreads in $\text{PG}(3, q)$, see e.g. [GS03a].

For vector space partitions of type $2^{m_2} 1^{m_1}$ of $\text{PG}(4, q)$ the packing condition in Equation (1) only implies $m_2 = q^3 + q - j$ and $m_1 = 1 + j(q+1)$ for $0 \leq j \leq q^3 + q$. However, the $1 + j(q+1)$ points that are not covered by the lines have to correspond to a (projective) q -divisible linear codes over \mathbb{F}_q of effective length $1 + j(q+1)$. Using the characterization result for the possible length of q^r -divisible codes over \mathbb{F}_q from [KK20] we can conclude $j \geq q-1$. Using this and the packing and the dimension condition in equations (1) and (2), we conclude:

Proposition 24. *The possible types of vector space partitions of $\text{PG}(4, q)$ are given by $4^1 1^{q^4}$, $3^1 2^{q^3-j} 1^{j(q+1)}$ for $0 \leq j \leq q^3$, and $2^{q^3+1-j} 1^{q^2+j(q+1)}$ for $0 \leq j \leq q^3 + 1$.*

All of the mentioned types can indeed be attained. Choosing an arbitrary solid in the ambient space leaves q^4 uncovered points. A lifted MRD code gives rise to a vector space partition \mathcal{P} of type $3^1 2^{q^3}$. In \mathcal{P} we can either replace j lines by their contained points or replace the plane by a line and q^2 points to obtain a vector space partition \mathcal{P}' of type $2^{q^3+1-j} 1^{q^2}$. In \mathcal{P}' we can then replace j lines by their contained points.

Using the same methods one can easily characterize all feasible types of vector space partitions in $\text{PG}(5, q)$ that do not contain a plane:

- $5^1 1^{q^5}$;
- $4^1 2^{q^4-j} 1^{j(q+1)}$ for $0 \leq j \leq q^4$;
- $2^{q^4+q^2+1-j} 1^{j(q+1)}$ for $0 \leq j \leq q^4 + q^2 + 1$.

A plane spread in $\text{PG}(5, q)$ is a vector space partition of type 3^{q^3+1} . From there we can easily obtain vector space partitions of type $3^{q^3+1-j} 2^{j-i} 1^{i(q+1)}$ for all $0 \leq j \leq q^3 + 1$ and all $0 \leq i \leq j$. However, also vector space partitions of other types do exist. With increasing dimension of the ambient space the problem of the classification of all feasible types of vector space partitions gets harder and harder. We would like to mention that in $\text{PG}(7, 3)$ the maximum number $A_3(8, 6; 3)$ of pairwise disjoint planes is unknown. The currently best known bounds are $244 \leq A_3(8, 6; 3) \leq 248$, see e.g. [HKK18]. If 248 such pairwise disjoint planes exist in

$\text{PG}(7, 3)$, then the 56 uncovered points have to form a so-called *Hill cap* [Hil78] corresponding to a two-weight code. Since the support of this object does not contain a line, there is e.g. no vector space partition of type $3^{248}2^11^{52}$ in $\text{PG}(7, 3)$.

References

- [BBK21] Iliya Bouyukliev, Stefka Bouyuklieva, and Sascha Kurz. Computer classification of linear codes. *IEEE Transactions on Information Theory*, 67(12):7807–7814, 2021.
- [BFIK22] John Bamberg, Yuval Filmus, Ferdinand Ihringer, and Sascha Kurz. Affine vector space partitions. *arXiv preprint 2211.16561*, 2022.
- [BFWW06] Iliya Bouyukliev, Veerle Fack, Wolfgang Willems, and Joost Winne. Projective two-weight codes with small parameters and their corresponding graphs. *Designs, Codes and Cryptography*, 41(1):59–78, 2006.
- [Bou07] Iliya Bouyukliev. What is Q-extension? *Serdica Journal of Computing*, 1(2):115–130, 2007.
- [DBDMS19] Jan De Beule, Jeroen Demeyer, Sam Mattheus, and Péter Sziklai. On the cylinder conjecture. *Designs, Codes and Cryptography*, 87(4):879–893, 2019.
- [EZHS⁺10] Saad Ibrahim El-Zanati, Olof Heden, George Francis Seelinger, Papa Amar Sissokho, Lawrence Edward Spence, and Charles Vanden Eynden. Partitions of the 8-dimensional vector space over $\text{GF}(2)$. *Journal of Combinatorial Designs*, 18(6):462–474, 2010.
- [EZJS⁺10] Saad El-Zanati, Heather Jordon, George Seelinger, Papa Sissokho, and Lawrence Spence. The maximum size of a partial 3-spread in a finite vector space over $\text{GF}(2)$. *Designs, Codes and Cryptography*, 54:101–107, 2010.
- [EZSS⁺09] Saad Ibrahim El-Zanati, George Francis Seelinger, Papa Amar Sissokho, Lawrence Edward Spence, and Charles Vanden Eynden. On partitions of finite vector spaces of low dimension over $\text{GF}(2)$. *Discrete Mathematics*, 309(14):4727–4735, 2009.
- [EZSS⁺11] Saad Ibrahim El-Zanati, George Francis Seelinger, Papa Amar Sissokho, Lawrence Edward Spence, and Charles Vanden Eynden. On λ -fold partitions of finite vector spaces and duality. *Discrete Mathematics*, 311(4):307–318, 2011.
- [GS03a] András Gács and Tamás Szőnyi. On maximal partial spreads in $\text{PG}(n, q)$. *Designs, Codes and Cryptography*, 29:123–129, 2003.
- [GS03b] Patrick Govaerts and Leo Storme. On a particular class of minihypers and its applications. I. The result for general q . *Designs, Codes and Cryptography*, 28(1):51–63, 2003.
- [Hed86] Olof Heden. Partitions of finite abelian groups. *European Journal of Combinatorics*, 7(1):11–25, 1986.
- [Hed09] Olof Heden. On the length of the tail of a vector space partition. *Discrete Mathematics*, 309(21):6169–6180, 2009.
- [Hed12] Olof Heden. A survey of the different types of vector space partitions. *Discrete Mathematics, Algorithms and Applications*, 4(01):1250001, 2012.
- [HHK⁺17] Daniel Heinlein, Thomas Honold, Michael Kiermaier, Sascha Kurz, and Alfred Wassermann. Projective divisible binary codes. In *The Tenth International Workshop on Coding and Cryptography 2017 : WCC Proceedings*. IEEE Information Theory Society, Saint-Petersburg, September 2017.

- [HHKK19] Daniel Heinlein, Thomas Honold, Michael Kiermaier, and Sascha Kurz. Generalized vector space partitions. *Australasian Journal of Combinatorics*, 73(1):162–178, 2019.
- [Hil78] Raymond Hill. Caps and codes. *Discrete Mathematics*, 22(2):111–137, 1978.
- [HKK18] Thomas Honold, Michael Kiermaier, and Sascha Kurz. Partial spreads and vector space partitions. In *Network Coding and Subspace Designs*, pages 131–170. Springer, 2018.
- [HKKW19] Thomas Honold, Michael Kiermaier, Sascha Kurz, and Alfred Wassermann. The lengths of projective triply-even binary codes. *IEEE Transactions on Information Theory*, 66(5):2713–2716, 2019.
- [HLNS13] Olof Heden, Juliane Lehmann, Esmeralda Năstase, and Papa Sissokho. The supertail of a subspace partition. *Designs, Codes and Cryptography*, 69(3):305–316, 2013.
- [KK08] Axel Kohnert and Sascha Kurz. Construction of large constant dimension codes with a prescribed minimum distance. *Mathematical Methods in Computer Science: Essays in Memory of Thomas Beth*, pages 31–42, 2008.
- [KK20] Michael Kiermaier and Sascha Kurz. On the lengths of divisible codes. *IEEE Transactions on Information Theory*, 66(7):4051–4060, 2020.
- [KM76] Earl S. Kramer and Dale M. Mesner. t -designs on hypergraphs. *Discrete Mathematics*, 15(3):263–296, 1976.
- [KM21] Sascha Kurz and Sam Mattheus. A generalization of the cylinder conjecture for divisible codes. *IEEE Transactions on Information Theory*, 68(4):2281–2289, 2021.
- [Kur21] Sascha Kurz. Divisible codes. *arXiv preprint 2112.11763*, 2021.
- [LH12] Juliane Lehmann and Olof Heden. Some necessary conditions for vector space partitions. *Discrete Mathematics*, 312(2):351–361, 2012.
- [Lun99] Guglielmo Lunardon. Normal spreads. *Geometriae Dedicata*, 75:245–261, 1999.
- [NS18] Esmeralda L. Năstase and Papa A. Sissokho. The complete characterization of the minimum size supertail in a subspace partition. *Linear Algebra and its Applications*, 559:172–180, 2018.
- [SSW19] John Sheekey, Kai-Uwe Schmidt, and Arne Winterhof. MRD codes: constructions and connections. In *Combinatorics and finite fields: Difference sets, polynomials, pseudorandomness and applications*, volume 23, pages 255–286. de Gruyter, 2019.

sascha@sascha-ThinkPad

A A more explicit variant of the main theorem

During the paper we have concluded several forbidden supertails. For the ease of the reader we summarize those that have been necessary in the classification of the feasible types of vector space partitions of $\text{PG}(7, 2)$:

Proposition 25. *No vector space partition of $\text{PG}(v - 1, 2)$ exists if it has a supertail of one of the following types:*

- 1^i for $1 \leq i \leq 2$;
- $2^0 1^4$;
- $2^{2-i} 1^{2+3i}$ for $0 \leq i \leq 1$;

- $2^{3-i}1^{3i}$ for all $0 \leq i \leq 3$;
- $2^{4-i}1^{3i}$ for all $0 \leq i \leq 4$;
- $2^{4-i}1^{1+3i}$ for all $0 \leq i \leq 4$;
- 2^31^5 ;
- 2^41^5 ;
- $3^32^{3-i}1^{3i}$ for $0 \leq i \leq 3$;
- $3^22^{5-i}1^{1+3i}$ for all $0 \leq i \leq 4$;
- $3^12^61^5$;
- $3^12^{11}1^5$.

With this we can reformulate Theorem 22 as follows:

Theorem 26. *Let \mathcal{P} be a vector space partition of $\text{PG}(7, 2)$ of type $7^{m_7}1^{m_1}$ satisfying the packing condition in Equation (1) and the dimension condition in Equation (2) as well as the special tail conditions from Proposition 25. Then, the type of \mathcal{P} is not contained in the following exhaustive list:*

- $4^{13}3^62^{6-i}1^{3i}$ for $0 \leq i \leq 6$;
- $4^{13}3^52^{7-i}1^{4+3i}$ for $0 \leq i \leq 7$;
- $4^{13}3^42^91^5$;
- $4^{12}3^82^{5-i}1^{4+3i}$ for $2 \leq i \leq 5$;
- $4^{11}3^{10}2^51^5$;
- $4^43^{25}2^51^5$.

A more explicit and extensive variant of our main theorem is given as follows:

Proposition 27. *Let \mathcal{P} be a vector space partition of $\text{PG}(7, 2)$, then \mathcal{P} has one of the following types:*

- 7^21^{128} ;
- $6^22^{64-i}1^{3i}$, where $0 \leq i \leq 64$;
- 5^13^{32} ;
- $5^13^{31}2^{1-i}1^{4+3i}$, where $0 \leq i \leq 1$;
- $5^13^{29}2^{7-i}1^{3i}$, where $0 \leq i \leq 7$;
- $5^13^{28}2^{8-i}1^{4+3i}$, where $0 \leq i \leq 8$;
- $5^13^{27}2^{10-i}1^{5+3i}$, where $0 \leq i \leq 10$;
- $5^13^{26}2^{14-i}1^{3i}$, where $0 \leq i \leq 14$;
- $5^13^{25}2^{15-i}1^{4+3i}$, where $0 \leq i \leq 15$;
- $5^13^{24}2^{17-i}1^{5+3i}$, where $0 \leq i \leq 17$;
- $5^13^{23}2^{21-i}1^{3i}$, where $0 \leq i \leq 21$;

- $5^1 3^{22} 2^{22-i} 1^{4+3i}$, where $0 \leq i \leq 22$;
- $5^1 3^{21} 2^{24-i} 1^{5+3i}$, where $0 \leq i \leq 24$;
- $5^1 3^{20} 2^{28-i} 1^{3i}$, where $0 \leq i \leq 28$;
- $5^1 3^{19} 2^{29-i} 1^{4+3i}$, where $0 \leq i \leq 29$;
- $5^1 3^{18} 2^{31-i} 1^{5+3i}$, where $0 \leq i \leq 31$;
- $5^1 3^{17} 2^{35-i} 1^{3i}$, where $0 \leq i \leq 35$;
- $5^1 3^{16} 2^{36-i} 1^{4+3i}$, where $0 \leq i \leq 36$;
- $5^1 3^{15} 2^{38-i} 1^{5+3i}$, where $0 \leq i \leq 38$;
- $5^1 3^{14} 2^{42-i} 1^{3i}$, where $0 \leq i \leq 42$;
- $5^1 3^{13} 2^{43-i} 1^{4+3i}$, where $0 \leq i \leq 43$;
- $5^1 3^{12} 2^{45-i} 1^{5+3i}$, where $0 \leq i \leq 45$;
- $5^1 3^{11} 2^{49-i} 1^{3i}$, where $0 \leq i \leq 49$;
- $5^1 3^{10} 2^{50-i} 1^{4+3i}$, where $0 \leq i \leq 50$;
- $5^1 3^9 2^{52-i} 1^{5+3i}$, where $0 \leq i \leq 52$;
- $5^1 3^8 2^{56-i} 1^{3i}$, where $0 \leq i \leq 56$;
- $5^1 3^7 2^{57-i} 1^{4+3i}$, where $0 \leq i \leq 57$;
- $5^1 3^6 2^{59-i} 1^{5+3i}$, where $0 \leq i \leq 59$;
- $5^1 3^5 2^{63-i} 1^{3i}$, where $0 \leq i \leq 63$;
- $5^1 3^4 2^{64-i} 1^{4+3i}$, where $0 \leq i \leq 64$;
- $5^1 3^3 2^{66-i} 1^{5+3i}$, where $0 \leq i \leq 66$;
- $5^1 3^2 2^{70-i} 1^{3i}$, where $0 \leq i \leq 70$;
- $5^1 3^1 2^{71-i} 1^{4+3i}$, where $0 \leq i \leq 71$;
- $5^1 2^{73-i} 1^{5+3i}$, where $0 \leq i \leq 73$;
- 4^{17} ;
- $4^{16} 3^1 1^8$;
- $4^{16} 2^{5-i} 1^{3i}$, where $0 \leq i \leq 5$;
- $4^{15} 3^2 1^{16}$;
- $4^{15} 3^1 2^{5-i} 1^{8+3i}$, where $0 \leq i \leq 5$;
- $4^{15} 2^{10-i} 1^{3i}$, where $0 \leq i \leq 10$;
- $4^{14} 3^3 2^{8-i} 1^{3i}$, where $0 \leq i \leq 8$;
- $4^{14} 3^2 2^{9-i} 1^{4+3i}$, where $0 \leq i \leq 9$;
- $4^{14} 3^1 2^{10-i} 1^{8+3i}$, where $0 \leq i \leq 10$;

- $4^{14}3^02^{15-i}1^{3i}$, where $0 \leq i \leq 15$;
- $4^{13}3^42^{8-i}1^{8+3i}$, where $0 \leq i \leq 8$;
- $4^{13}3^32^{13-i}1^{3i}$, where $0 \leq i \leq 13$;
- $4^{13}3^22^{14-i}1^{4+3i}$, where $0 \leq i \leq 14$;
- $4^{13}3^12^{16-i}1^{5+3i}$ where $0 \leq i \leq 16$;
- $4^{13}3^02^{20-i}1^{3i}$, where $0 \leq i \leq 20$;
- $4^{12}3^82^{1-i}1^{16+3i}$, where $0 \leq i \leq 1$;
- $4^{12}3^72^{7-i}1^{5+3i}$, where $0 \leq i \leq 7$;
- $4^{12}3^62^{11-i}1^{3i}$, where $0 \leq i \leq 11$;
- $4^{12}3^52^{12-i}1^{4+3i}$, where $0 \leq i \leq 11$;
- $4^{12}3^42^{14-i}1^{5+3i}$, where $0 \leq i \leq 14$;
- $4^{12}3^32^{18-i}1^{3i}$, where $0 \leq i \leq 18$;
- $4^{12}3^22^{19-i}1^{4+3i}$, where $0 \leq i \leq 19$;
- $4^{12}3^12^{21-i}1^{5+3i}$, where $0 \leq i \leq 21$;
- $4^{12}2^{25-i}1^{3i}$, where $0 \leq i \leq 25$;
- $4^{11}3^102^{4-i}1^{8+3i}$, where $0 \leq i \leq 4$;
- $4^{11}3^92^{9-i}1^{3i}$, where $0 \leq i \leq 9$;
- $4^{11}3^82^{10-i}1^{4+3i}$, where $0 \leq i \leq 10$;
- $4^{11}3^72^{12-i}1^{5+3i}$, where $0 \leq i \leq 12$;
- $4^{11}3^62^{16-i}1^{0}3i$, where $0 \leq i \leq 16$;
- $4^{11}3^52^{17-i}1^{4+3i}$, where $0 \leq i \leq 17$;
- $4^{11}3^42^{19-i}1^{5+3i}$, where $0 \leq i \leq 19$;
- $4^{11}3^32^{23-i}1^{3i}$, where $0 \leq i \leq 23$;
- $4^{11}3^22^{24-i}1^{4+3i}$, where $0 \leq i \leq 24$;
- $4^{11}3^12^{26-i}1^{5+3i}$, where $0 \leq i \leq 26$;
- $4^{11}2^{30-i}1^{3i}$, where $0 \leq i \leq 30$;
- $4^{10}3^{15}$;
- $4^{10}3^{14}2^{1-i}1^{4+3i}$, where $0 \leq i \leq 1$;
- $4^{10}3^{13}2^{2-i}1^{8+3i}$, where $0 \leq i \leq 2$;
- $4^{10}3^{12}2^{7-i}1^{3i}$, where $0 \leq i \leq 7$;
- $4^{10}3^{11}2^{8-i}1^{4+3i}$, where $0 \leq i \leq 8$;
- $4^{10}3^{10}2^{10-i}1^{5+3i}$, where $0 \leq i \leq 10$;

- $4^{10}3^92^{14-i}1^{3i}$, where $0 \leq i \leq 14$;
- $4^{10}3^82^{15-i}1^{4+3i}$, where $0 \leq i \leq 15$;
- $4^{10}3^72^{17-i}1^{5+3i}$, where $0 \leq i \leq 17$;
- $4^{10}3^62^{21-i}1^{3i}$, where $0 \leq i \leq 21$;
- $4^{10}3^52^{22-i}1^{4+3i}$, where $0 \leq i \leq 22$;
- $4^{10}3^42^{24-i}1^{5+3i}$, where $0 \leq i \leq 24$;
- $4^{10}3^32^{28-i}1^{3i}$, where $0 \leq i \leq 28$;
- $4^{10}3^22^{29-i}1^{4+3i}$, where $0 \leq i \leq 29$;
- $4^{10}3^12^{31-i}1^{5+3i}$, where $0 \leq i \leq 31$;
- $4^{10}2^{35-i}1^{3i}$, where $0 \leq i \leq 35$;
- $4^93^{16}1^8$;
- $4^93^{15}2^{5-i}1^{3i}$, where $0 \leq i \leq 5$;
- $4^93^{14}2^{6-i}1^{4+3i}$, where $0 \leq i \leq 6$;
- $4^93^{13}2^{8-i}1^{5+3i}$, where $0 \leq i \leq 8$;
- $4^93^{12}2^{12-i}1^{3i}$, where $0 \leq i \leq 12$;
- $4^93^{11}2^{13-i}1^{4+3i}$, where $0 \leq i \leq 13$;
- $4^93^{10}2^{15-i}1^{5+3i}$, where $0 \leq i \leq 15$;
- $4^93^92^{19-i}1^{3i}$, where $0 \leq i \leq 19$;
- $4^93^82^{20-i}1^{4+3i}$, where $0 \leq i \leq 20$;
- $4^93^72^{22-i}1^{5+3i}$, where $0 \leq i \leq 22$;
- $4^93^62^{26-i}1^{3i}$, where $0 \leq i \leq 26$;
- $4^93^52^{27-i}1^{4+3i}$, where $0 \leq i \leq 27$;
- $4^93^42^{29-i}1^{5+3i}$, where $0 \leq i \leq 29$;
- $4^93^32^{33-i}1^{3i}$, where $0 \leq i \leq 33$;
- $4^93^22^{34-i}1^{4+3i}$, where $0 \leq i \leq 34$;
- $4^93^12^{36-i}1^{5+3i}$, where $0 \leq i \leq 36$;
- $4^92^{40-i}1^{3i}$, where $0 \leq i \leq 40$;
- $4^83^{17}2^41^{4+3i}$, where $0 \leq i \leq 4$;
- $4^83^{16}2^61^{5+3i}$, where $0 \leq i \leq 6$;
- $4^83^{15}2^{10-i}1^{3i}$, where $0 \leq i \leq 10$;
- $4^83^{14}2^{11-i}1^{4+3i}$, where $0 \leq i \leq 11$;
- $4^83^{13}2^{13-i}1^{5+3i}$, where $0 \leq i \leq 13$;

- $4^8 3^{12} 2^{17-i} 1^{3i}$, where $0 \leq i \leq 17$;
- $4^8 3^{11} 2^{18-i} 1^{4+3i}$, where $0 \leq i \leq 18$;
- $4^8 3^{10} 2^{20-i} 1^{5+3i}$, where $0 \leq i \leq 20$;
- $4^8 3^9 2^{24-i} 1^{3i}$, where $0 \leq i \leq 24$;
- $4^8 3^8 2^{25-i} 1^{4+3i}$, where $0 \leq i \leq 25$;
- $4^8 3^7 2^{27-i} 1^{5+3i}$, where $0 \leq i \leq 27$;
- $4^8 3^6 2^{31-i} 1^{3i}$, where $0 \leq i \leq 31$;
- $4^8 3^5 2^{32-i} 1^{4+3i}$, where $0 \leq i \leq 32$;
- $4^8 3^4 2^{34-i} 1^{5+3i}$, where $0 \leq i \leq 34$;
- $4^8 3^3 2^{38-i} 1^{3i}$, where $0 \leq i \leq 38$;
- $4^8 3^2 2^{39-i} 1^{4+3i}$, where $0 \leq i \leq 39$;
- $4^8 3^1 2^{41-i} 1^{5+3i}$, where $0 \leq i \leq 41$;
- $4^8 3^0 2^{45-i} 1^{3i}$, where $0 \leq i \leq 45$;
- $4^7 3^{19} 2^{3-i} 1^{8+3i}$, where $0 \leq i \leq 3$;
- $4^7 3^{18} 2^{8-i} 1^{3i}$, where $0 \leq i \leq 8$;
- $4^7 3^{17} 2^{9-i} 1^{4+3i}$, where $0 \leq i \leq 9$;
- $4^7 3^{16} 2^{11-i} 1^{5+3i}$, where $0 \leq i \leq 11$;
- $4^7 3^{15} 2^{15-i} 1^{0+3i}$, where $0 \leq i \leq 15$;
- $4^7 3^{14} 2^{16-i} 1^{4+3i}$, where $0 \leq i \leq 16$;
- $4^7 3^{13} 2^{18-i} 1^{5+3i}$, where $0 \leq i \leq 18$;
- $4^7 3^{12} 2^{22-i} 1^{3i}$, where $0 \leq i \leq 22$;
- $4^7 3^{11} 2^{23-i} 1^{4+3i}$, where $0 \leq i \leq 23$;
- $4^7 3^{10} 2^{25-i} 1^{5+3i}$, where $0 \leq i \leq 25$;
- $4^7 3^9 2^{29-i} 1^{3i}$, where $0 \leq i \leq 29$;
- $4^7 3^8 2^{30-i} 1^{4+3i}$, where $0 \leq i \leq 30$;
- $4^7 3^7 2^{32-i} 1^{5+3i}$, where $0 \leq i \leq 32$;
- $4^7 3^6 2^{36-i} 1^{3i}$, where $0 \leq i \leq 36$;
- $4^7 3^5 2^{37-i} 1^{4+3i}$, where $0 \leq i \leq 37$;
- $4^7 3^4 2^{39-i} 1^{5+3i}$, where $0 \leq i \leq 39$;
- $4^7 3^3 2^{43-i} 1^{3i}$, where $0 \leq i \leq 43$;
- $4^7 3^2 2^{44-i} 1^{4+3i}$, where $0 \leq i \leq 44$;
- $4^7 3^1 2^{46-i} 1^{5+3i}$, where $0 \leq i \leq 46$;

- $4^7 2^{50-i} 1^{3i}$, where $0 \leq i \leq 50$;
- $4^6 3^{21} 2^{6-i} 1^{3i}$, where $0 \leq i \leq 6$;
- $4^6 3^{20} 2^{7-i} 1^{4+3i}$, where $0 \leq i \leq 7$;
- $4^6 3^{19} 2^{9-i} 1^{5+3i}$, where $0 \leq i \leq 9$;
- $4^6 3^{18} 2^{13-i} 1^{3i}$, where $0 \leq i \leq 13$;
- $4^6 3^{17} 2^{14-i} 1^{4+3i}$, where $0 \leq i \leq 14$;
- $4^6 3^{16} 2^{16-i} 1^{5+3i}$, where $0 \leq i \leq 16$;
- $4^6 3^{15} 2^{20-i} 1^{3i}$, where $0 \leq i \leq 20$;
- $4^6 3^{14} 2^{21-i} 1^{4+3i}$, where $0 \leq i \leq 21$;
- $4^6 3^{13} 2^{23-i} 1^{5+3i}$, where $0 \leq i \leq 23$;
- $4^6 3^{12} 2^{27-i} 1^{3i}$, where $0 \leq i \leq 27$;
- $4^6 3^{11} 2^{28-i} 1^{4+3i}$, where $0 \leq i \leq 28$;
- $4^6 3^{10} 2^{30-i} 1^{5+3i}$, where $0 \leq i \leq 30$;
- $4^6 3^9 2^{34-i} 1^{3i}$, where $0 \leq i \leq 34$;
- $4^6 3^8 2^{35-i} 1^{4+3i}$, where $0 \leq i \leq 35$;
- $4^6 3^7 2^{37-i} 1^{5+3i}$, where $0 \leq i \leq 37$;
- $4^6 3^6 2^{41-i} 1^{3i}$, where $0 \leq i \leq 41$;
- $4^6 3^5 2^{42-i} 1^{4+3i}$, where $0 \leq i \leq 42$;
- $4^6 3^4 2^{44-i} 1^{5+3i}$, where $0 \leq i \leq 44$;
- $4^6 3^3 2^{48-i} 1^{3i}$, where $0 \leq i \leq 48$;
- $4^6 3^2 2^{49-i} 1^{4+3i}$, where $0 \leq i \leq 49$;
- $4^6 3^1 2^{51-i} 1^{5+3i}$, where $0 \leq i \leq 51$;
- $4^6 3^0 2^{55-i} 1^{3i}$, where $0 \leq i \leq 55$;
- $4^5 3^{23} 2^{5-i} 1^{4+3i}$, where $0 \leq i \leq 5$;
- $4^5 3^{22} 2^{7-i} 1^{5+3i}$, where $0 \leq i \leq 7$;
- $4^5 3^{21} 2^{11-i} 1^{3i}$, where $0 \leq i \leq 11$;
- $4^5 3^{20} 2^{12-i} 1^{4+3i}$, where $0 \leq i \leq 12$;
- $4^5 3^{19} 2^{14-i} 1^{5+3i}$, where $0 \leq i \leq 14$;
- $4^5 3^{18} 2^{18-i} 1^{3i}$, where $0 \leq i \leq 18$;
- $4^5 3^{17} 2^{19-i} 1^{4+3i}$, where $0 \leq i \leq 19$;
- $4^5 3^{16} 2^{21-i} 1^{5+3i}$, where $0 \leq i \leq 21$;
- $4^5 3^{15} 2^{25-i} 1^{3i}$, where $0 \leq i \leq 25$;

- $4^5 3^{14} 2^{26-i} 1^{4+3i}$, where $0 \leq i \leq 26$;
- $4^5 3^{13} 2^{28-i} 1^{5+3i}$, where $0 \leq i \leq 28$;
- $4^5 3^{12} 2^{32-i} 1^{3i}$, where $0 \leq i \leq 32$;
- $4^5 3^{11} 2^{33-i} 1^{4+3i}$, where $0 \leq i \leq 33$;
- $4^5 3^{10} 2^{35-i} 1^{5+3i}$, where $0 \leq i \leq 35$;
- $4^5 3^9 2^{39-i} 1^{3i}$, where $0 \leq i \leq 39$;
- $4^5 3^8 2^{40-i} 1^{4+3i}$, where $0 \leq i \leq 40$;
- $4^5 3^7 2^{42-i} 1^{5+3i}$, where $0 \leq i \leq 42$;
- $4^5 3^6 2^{46-i} 1^{3i}$, where $0 \leq i \leq 46$;
- $4^5 3^5 2^{47-i} 1^{4+3i}$, where $0 \leq i \leq 47$;
- $4^5 3^4 2^{49-i} 1^{5+3i}$, where $0 \leq i \leq 49$;
- $4^5 3^3 2^{53-i} 1^{3i}$, where $0 \leq i \leq 53$;
- $4^5 3^2 2^{54-i} 1^{4+3i}$, where $0 \leq i \leq 54$;
- $4^5 3^1 2^{56-i} 1^{5+3i}$, where $0 \leq i \leq 56$;
- $4^5 2^{60-i} 1^{3i}$, where $0 \leq i \leq 60$;
- $4^4 3^{25} 2^{4-i} 1^{8+3i}$, where $0 \leq i \leq 4$;
- $4^4 3^{24} 2^{9-i} 1^{3i}$, where $0 \leq i \leq 9$;
- $4^4 3^{23} 2^{10-i} 1^{4+3i}$, where $0 \leq i \leq 10$;
- $4^4 3^{22} 2^{12-i} 1^{5+3i}$, where $0 \leq i \leq 12$;
- $4^4 3^{21} 2^{16-i} 1^{3i}$, where $0 \leq i \leq 16$;
- $4^4 3^{20} 2^{17-i} 1^{4+3i}$, where $0 \leq i \leq 17$;
- $4^4 3^{19} 2^{19-i} 1^{5+3i}$, where $0 \leq i \leq 19$;
- $4^4 3^{18} 2^{23-i} 1^{3i}$, where $0 \leq i \leq 23$;
- $4^4 3^{17} 2^{24-i} 1^{4+3i}$, where $0 \leq i \leq 24$;
- $4^4 3^{16} 2^{26-i} 1^{5+3i}$, where $0 \leq i \leq 26$;
- $4^4 3^{15} 2^{30-i} 1^{3i}$, where $0 \leq i \leq 30$;
- $4^4 3^{14} 2^{31-i} 1^{4+3i}$, where $0 \leq i \leq 31$;
- $4^4 3^{13} 2^{33-i} 1^{5+3i}$, where $0 \leq i \leq 33$;
- $4^4 3^{12} 2^{37-i} 1^{3i}$, where $0 \leq i \leq 37$;
- $4^4 3^{11} 2^{38-i} 1^{4+3i}$, where $0 \leq i \leq 38$;
- $4^4 3^{10} 2^{40-i} 1^{5+3i}$, where $0 \leq i \leq 40$;
- $4^4 3^9 2^{44-i} 1^{3i}$, where $0 \leq i \leq 44$;

- $4^4 3^8 2^{45-i} 1^{4+3i}$, where $0 \leq i \leq 45$;
- $4^4 3^7 2^{47-i} 1^{5+3i}$, where $0 \leq i \leq 47$;
- $4^4 3^6 2^{51-i} 1^{3i}$, where $0 \leq i \leq 51$;
- $4^4 3^5 2^{52-i} 1^{4+3i}$, where $0 \leq i \leq 52$;
- $4^4 3^4 2^{54-i} 1^{5+3i}$, where $0 \leq i \leq 54$;
- $4^4 3^3 2^{58-i} 1^{3i}$, where $0 \leq i \leq 58$;
- $4^4 3^2 2^{59-i} 1^{4+3i}$, where $0 \leq i \leq 59$;
- $4^4 3^1 2^{61-i} 1^{5+3i}$, where $0 \leq i \leq 61$;
- $4^4 3^0 2^{65-i} 1^{3i}$, where $0 \leq i \leq 65$;
- $4^3 3^{30}$;
- $4^3 3^{29} 2^{1-i} 1^4 + 3i$, where $0 \leq i \leq 1$;
- $4^3 3^{28} 2^{2-i} 1^{8+3i}$, where $0 \leq i \leq 2$;
- $4^3 3^{27} 2^{7-i} 1^{3i}$, where $0 \leq i \leq 7$;
- $4^3 3^{26} 2^{8-i} 1^{4+3i}$, where $0 \leq i \leq 8$;
- $4^3 3^{25} 2^{10-i} 1^{5+3i}$, where $0 \leq i \leq 10$;
- $4^3 3^{24} 2^{14-i} 1^{3i}$, where $0 \leq i \leq 14$;
- $4^3 3^{23} 2^{15-i} 1^{4+3i}$, where $0 \leq i \leq 15$;
- $4^3 3^{22} 2^{17-i} 1^{5+3i}$, where $0 \leq i \leq 17$;
- $4^3 3^{21} 2^{21-i} 1^{3i}$, where $0 \leq i \leq 21$;
- $4^3 3^{20} 2^{22-i} 1^{4+3i}$, where $0 \leq i \leq 22$;
- $4^3 3^{19} 2^{24-i} 1^{5+3i}$, where $0 \leq i \leq 24$;
- $4^3 3^{18} 2^{28-i} 1^{3i}$, where $0 \leq i \leq 2$;
- $4^3 3^{17} 2^{29-i} 1^{4+3i}$, where $0 \leq i \leq 29$;
- $4^3 3^{16} 2^{31-i} 1^{5+3i}$, where $0 \leq i \leq 31$;
- $4^3 3^{15} 2^{35-i} 1^{3i}$, where $0 \leq i \leq 35$;
- $4^3 3^{14} 2^{36-i} 1^{4+3i}$, where $0 \leq i \leq 36$;
- $4^3 3^{13} 2^{38-i} 1^{5+3i}$, where $0 \leq i \leq 38$;
- $4^3 3^{12} 2^{42-i} 1^{3i}$, where $0 \leq i \leq 42$;
- $4^3 3^{11} 2^{43-i} 1^{4+3i}$, where $0 \leq i \leq 43$;
- $4^3 3^{10} 2^{45-i} 1^{5+3i}$, where $0 \leq i \leq 45$;
- $4^3 3^9 2^{49-i} 1^{3i}$, where $0 \leq i \leq 49$;
- $4^3 3^8 2^{50-i} 1^{4+3i}$, where $0 \leq i \leq 50$;

- $4^3 3^7 2^{52-i} 1^{5+3i}$, where $0 \leq i \leq 52$;
- $4^3 3^6 2^{56-i} 1^{3i}$, where $0 \leq i \leq 56$;
- $4^3 3^5 2^{57-i} 1^{4+3i}$, where $0 \leq i \leq 57$;
- $4^3 3^4 2^{59-i} 1^{5+3i}$, where $0 \leq i \leq 59$;
- $4^3 3^3 2^{63-i} 1^{3i}$, where $0 \leq i \leq 63$;
- $4^3 3^2 2^{64-i} 1^{4+3i}$, where $0 \leq i \leq 64$;
- $4^3 3^1 2^{66-i} 1^{5+3i}$, where $0 \leq i \leq 66$;
- $4^3 3^0 2^{70-i} 1^{3i}$, where $0 \leq i \leq 70$;
- $4^2 3^{31} 1^8$;
- $4^2 3^{30} 2^{5-i} 1^{3i}$, where $0 \leq i \leq 5$;
- $4^2 3^{29} 2^{6-i} 1^{4+3i}$, where $0 \leq i \leq 6$;
- $4^2 3^{28} 2^{8-i} 1^{5+3i}$, where $0 \leq i \leq 8$;
- $4^2 3^{27} 2^{12-i} 1^{3i}$, where $0 \leq i \leq 12$;
- $4^2 3^{26} 2^{13-i} 1^{4+3i}$, where $0 \leq i \leq 13$;
- $4^2 3^{25} 2^{15-i} 1^{5+3i}$, where $0 \leq i \leq 15$;
- $4^2 3^{24} 2^{19-i} 1^{3i}$, where $0 \leq i \leq 19$;
- $4^2 3^{23} 2^{20-i} 1^{4+3i}$, where $0 \leq i \leq 20$;
- $4^2 3^{22} 2^{22-i} 1^{5+3i}$, where $0 \leq i \leq 22$;
- $4^2 3^{21} 2^{26-i} 1^{3i}$, where $0 \leq i \leq 26$;
- $4^2 3^{20} 2^{27-i} 1^{4+3i}$, where $0 \leq i \leq 27$;
- $4^2 3^{19} 2^{29-i} 1^{5+3i}$, where $0 \leq i \leq 29$;
- $4^2 3^{18} 2^{33-i} 1^{3i}$, where $0 \leq i \leq 33$;
- $4^2 3^{17} 2^{34-i} 1^{4+3i}$, where $0 \leq i \leq 34$;
- $4^2 3^{16} 2^{36-i} 1^{5+3i}$, where $0 \leq i \leq 36$;
- $4^2 3^{15} 2^{40-i} 1^{3i}$, where $0 \leq i \leq 40$;
- $4^2 3^{14} 2^{41-i} 1^{4+3i}$, where $0 \leq i \leq 41$;
- $4^2 3^{13} 2^{43-i} 1^{5+3i}$, where $0 \leq i \leq 43$;
- $4^2 3^{12} 2^{47-i} 1^{3i}$, where $0 \leq i \leq 47$;
- $4^2 3^{11} 2^{48-i} 1^{4+3i}$, where $0 \leq i \leq 48$;
- $4^2 3^{10} 2^{50-i} 1^{5+3i}$, where $0 \leq i \leq 50$;
- $4^2 3^9 2^{54-i} 1^{3i}$, where $0 \leq i \leq 54$;
- $4^2 3^8 2^{55-i} 1^{4+3i}$, where $0 \leq i \leq 55$;

- $4^2 3^7 2^{57-i} 1^{5+3i}$, where $0 \leq i \leq 57$;
- $4^2 3^6 2^{61-i} 1^{3i}$, where $0 \leq i \leq 61$;
- $4^2 3^5 2^{62-i} 1^{4+3i}$, where $0 \leq i \leq 62$;
- $4^2 3^4 2^{64-i} 1^{5+3i}$, where $0 \leq i \leq 64$;
- $4^2 3^3 2^{68-i} 1^{3i}$, where $0 \leq i \leq 68$;
- $4^2 3^2 2^{69-i} 1^{4+3i}$, where $0 \leq i \leq 69$;
- $4^2 3^1 2^{71-i} 1^{5+3i}$, where $0 \leq i \leq 71$;
- $4^2 3^0 2^{75-i} 1^{3i}$, where $0 \leq i \leq 75$;
- $4^1 3^{32} 2^{4-i} 1^{4+3i}$, where $0 \leq i \leq 4$;
- $4^1 3^{31} 2^{6-i} 1^{5+3i}$, where $0 \leq i \leq 6$;
- $4^1 3^{30} 2^{10-i} 1^{3i}$, where $0 \leq i \leq 10$;
- $4^1 3^{29} 2^{11-i} 1^{4+3i}$, where $0 \leq i \leq 11$;
- $4^1 3^{28} 2^{13-i} 1^{5+3i}$, where $0 \leq i \leq 13$;
- $4^1 3^{27} 2^{17-i} 1^{3i}$, where $0 \leq i \leq 17$;
- $4^1 3^{26} 2^{18-i} 1^{4+3i}$, where $0 \leq i \leq 18$;
- $4^1 3^{25} 2^{20-i} 1^{5+3i}$, where $0 \leq i \leq 20$;
- $4^1 3^{24} 2^{24-i} 1^{3i}$, where $0 \leq i \leq 24$;
- $4^1 3^{23} 2^{25-i} 1^{4+3i}$, where $0 \leq i \leq 25$;
- $4^1 3^{22} 2^{27-i} 1^{5+3i}$, where $0 \leq i \leq 27$;
- $4^1 3^{21} 2^{31-i} 1^{3i}$, where $0 \leq i \leq 31$;
- $4^1 3^{20} 2^{32-i} 1^{4+3i}$, where $0 \leq i \leq 32$;
- $4^1 3^{19} 2^{34-i} 1^{5+3i}$, where $0 \leq i \leq 34$;
- $4^1 3^{18} 2^{38-i} 1^{3i}$, where $0 \leq i \leq 38$;
- $4^1 3^{17} 2^{39-i} 1^{4+3i}$, where $0 \leq i \leq 39$;
- $4^1 3^{16} 2^{41-i} 1^{5+3i}$, where $0 \leq i \leq 41$;
- $4^1 3^{15} 2^{45-i} 1^{3i}$, where $0 \leq i \leq 45$;
- $4^1 3^{14} 2^{46-i} 1^{4+3i}$, where $0 \leq i \leq 46$;
- $4^1 3^{13} 2^{48-i} 1^{5+3i}$, where $0 \leq i \leq 48$;
- $4^1 3^{12} 2^{52-i} 1^{3i}$, where $0 \leq i \leq 52$;
- $4^1 3^{11} 2^{53-i} 1^{4+3i}$, where $0 \leq i \leq 53$;
- $4^1 3^{10} 2^{55-i} 1^{5+3i}$, where $0 \leq i \leq 55$;
- $4^1 3^9 2^{59-i} 1^{3i}$, where $0 \leq i \leq 59$;

- $4^1 3^8 2^{60-i} 1^{4+3i}$, where $0 \leq i \leq 60$;
- $4^1 3^7 2^{62-i} 1^{5+3i}$, where $0 \leq i \leq 62$;
- $4^1 3^6 2^{66-i} 1^{3i}$, where $0 \leq i \leq 66$;
- $4^1 3^5 2^{67-i} 1^{4+3i}$, where $0 \leq i \leq 67$;
- $4^1 3^4 2^{69-i} 1^{5+3i}$, where $0 \leq i \leq 69$;
- $4^1 3^3 2^{73-i} 1^{3i}$, where $0 \leq i \leq 73$;
- $4^1 3^2 2^{74-i} 1^{4+3i}$, where $0 \leq i \leq 74$;
- $4^1 3^1 2^{76-i} 1^{5+3i}$, where $0 \leq i \leq 76$;
- $4^1 3^0 2^{80-i} 1^{3i}$, where $0 \leq i \leq 80$;
- $4^0 3^4 2^{3-i} 1^{8+3i}$, where $0 \leq i \leq 3$;
- $4^0 3^3 2^{8-i} 1^{3i}$, where $0 \leq i \leq 8$;
- $4^0 3^3 2^{9-i} 1^{4+3i}$, where $0 \leq i \leq 9$;
- $4^0 3^3 1 2^{11-i} 1^{5+3i}$, where $0 \leq i \leq 11$;
- $4^0 3^3 2^{15-i} 1^{3i}$, where $0 \leq i \leq 15$;
- $4^0 3^2 2^{16-i} 1^{4+3i}$, where $0 \leq i \leq 16$;
- $4^0 3^2 2^{18-i} 1^{5+3i}$, where $0 \leq i \leq 18$;
- $4^0 3^2 2^{22-i} 1^{3i}$, where $0 \leq i \leq 22$;
- $4^0 3^2 2^{23-i} 1^{4+3i}$, where $0 \leq i \leq 23$;
- $4^0 3^2 2^{25-i} 1^{5+3i}$, where $0 \leq i \leq 25$;
- $4^0 3^2 2^{29-i} 1^{3i}$, where $0 \leq i \leq 29$;
- $4^0 3^2 2^{30-i} 1^{4+3i}$, where $0 \leq i \leq 30$;
- $4^0 3^2 2^{32-i} 1^{5+3i}$, where $0 \leq i \leq 32$;
- $4^0 3^2 1 2^{36-i} 1^{3i}$, where $0 \leq i \leq 36$;
- $4^0 3^2 2^{37-i} 1^{4+3i}$, where $0 \leq i \leq 37$;
- $4^0 3^1 2^{39-i} 1^{5+3i}$, where $0 \leq i \leq 39$;
- $4^0 3^1 2^{43-i} 1^{3i}$, where $0 \leq i \leq 43$;
- $4^0 3^1 2^{44-i} 1^{4+3i}$, where $0 \leq i \leq 44$;
- $4^0 3^1 2^{46-i} 1^{5+3i}$, where $0 \leq i \leq 46$;
- $4^0 3^1 2^{50-i} 1^{3i}$, where $0 \leq i \leq 50$;
- $4^0 3^1 2^{51-i} 1^{4+3i}$, where $0 \leq i \leq 51$;
- $4^0 3^1 2^{53-i} 1^{5+3i}$, where $0 \leq i \leq 53$;
- $4^0 3^1 2^{57-i} 1^{3i}$, where $0 \leq i \leq 57$;

- $4^0 3^{11} 2^{58-i} 1^{4+3i}$, where $0 \leq i \leq 58$;
- $4^0 3^{10} 2^{60-i} 1^{5+3i}$, where $0 \leq i \leq 60$;
- $4^0 3^9 2^{64-i} 1^{3i}$, where $0 \leq i \leq 64$;
- $4^0 3^8 2^{65-i} 1^{4+3i}$, where $0 \leq i \leq 65$;
- $4^0 3^7 2^{67-i} 1^{5+3i}$, where $0 \leq i \leq 67$;
- $4^0 3^6 2^{71-i} 1^{3i}$, where $0 \leq i \leq 71$;
- $4^0 3^5 2^{72-i} 1^{4+3i}$, where $0 \leq i \leq 72$;
- $4^0 3^4 2^{74-i} 1^{5+3i}$, where $0 \leq i \leq 74$;
- $4^0 3^3 2^{78-i} 1^{3i}$, where $0 \leq i \leq 78$;
- $4^0 3^2 2^{79-i} 1^{4+3i}$, where $0 \leq i \leq 79$;
- $4^0 3^1 2^{81-i} 1^{5+3i}$, where $0 \leq i \leq 81$;
- $4^0 3^0 2^{85-i} 1^{3i}$, where $0 \leq i \leq 85$;